

## DAFTAR PUSTAKA

- Aban, S.M., Argente, F.A.T., Raguindin, R.S., Garcia, A.C., Ibarra, C.E. & De Vera, R.B., 2017. Length-weight relationships of the asian green mussel, *Perna viridis* (Linnaeus 1758) (Bivalvia: Mytilidae) population in Bolinao Bay, Pangasinan. *PSU Journal of Natural and Allied Sciences*, 1(1):1-6.
- Alnahdi, A., Leaniz, C.G.D., and King, A. J. (2016). Spatio-temporal variation in length-weight relationships and condition of the ribbonfish *Trichiurus lepturus* (Linnaeus, 1758): Implications for fisheries management. *Journal Plos One*, 11(8): 1-14.
- At, A., Erling, U. J. P., Angsa, K. O. T. A. L., & Ceh, A. (2018). Hubungan Panjang Dan Berat Serta Faktor Kondisi Kerang Bulu *Anadara antiquata* Di Ujung Perling, Kota Langsa Aceh. 2, 30-34.
- Aya, F. A., Hidaka, Y., & Kudo, I. (2014). Clearance rates and ingestion efficiency of the Japanese scallop *Patinopecten yessoensis*. *Plankton and Benthos Research*, 8(3), 134-140.
- Belcheva, N. N., Zakhartsev, M., Silina, A. V., Slinko, E. N., & Chelomin, V. P. (2006). Relationship between shell weight and cadmium content in whole digestive gland of the Japanese scallop *Patinopecten yessoensis* (Jay). *Marine Environmental Research*, 61(4), 396-409.
- Buban, I. C. R., Soliman, V. S., Bobiles, R. U., & Pulvinar, A. (2019). Morpho-biometric relationship, relative condition factor and meat yield of distant scallop *Bractechlamys vexillum* (Reeve, 1853) in Asid Gulf, Philippines. *Asian Fisheries Science*, 32(4), 147-153.
- Chauvaud, L., Patry, Y., Jolivet, A., Cam, E., Le Goff, C., Strand, Ø., Charrier, G., Thébault, J., Lazure, P., Gotthard, K., & Clavier, J. (2012). Variation in size and growth of the Great Scallop *Pecten maximus* along a latitudinal gradient. *PLoS ONE*, 7(5), 6-15.
- Ding, J., Zhao, L., Chang, Y., Zhao, W., Du, Z., & Hao, Z. (2015). Transcriptome sequencing and characterization of Japanese scallop *Patinopecten yessoensis* from different shell color lines. *PLoS ONE*, 10(2), 1-18.
- Dvoretzky, A. G., & Dvoretzky, V. G. (2022). Biological Aspects, Fisheries, and Aquaculture of Yesso Scallops in Russian Waters of the Sea of Japan. *Diversity*, 14(5).
- Fauzan, M., Bakti, D., Susetya, I. E., & Desrita. (2018). Growth and exploitation rate of *Anadara gubernaculum* (reeve, 1844) Arcidae

Family in Asahan Aquatic of North Sumatra. *IOP Conference Series: Earth and Environmental Science*, 122(1).

- Fisheriesaquaculture, F. A. O., & Paper, T. (2022). Hatchery-based seed production of the Japanese scallop, *Mizuhopecten yessoensis*. In *Hatchery-based seed production of the Japanese scallop, Mizuhopecten yessoensis*.
- Grant, J., Emerson, C. W., Mallet, A., & Carver, C. (2003). Growth advantages of ear hanging compared to cage culture for sea scallops, *Placopecten magellanicus* PII: S 0 0 4 4-8 4 8 6 ( 0 2 ) 0 0 3 6 0-5. In *Aquaculture* (Vol. 217).
- Hamada, T., Yamashita, N., Watanabe, T., & Natsume, S. (2001). Drilling position of the ear affects growth and mortality of scallop (*Patinopecten yessoensis*, Jay) in ear-hanging culture. *Aquaculture*, 193(3-4), 249-256.
- Hou, R., Bao, Z., Wang, S., Su, H., Li, Y., Du, H., Hu, J., Wang, S., & Hu, X. (2011). Transcriptome sequencing and De Novo analysis for Yesso Scallop (*Patinopecten yessoensis*) using 454 GS FLX. *PLoS ONE*, 6(6).
- Ichinokawa, M., Okamura, H., & Kurota, H. (2017). The status of Japanese fisheries relative to fisheries around the world. *ICES Journal of Marine Science*, 74(5), 1277-1287.
- Jiang, W., Li, J., Gao, Y., Mao, Y., Jiang, Z., Du, M., Zhang, Y., & Fang, J. (2016). Effects of temperature change on physiological and biochemical responses of Yesso scallop, *Patinopecten yessoensis*. *Aquaculture*, 451, 463-472.
- Kanamori, M., Baba, K., Natsuike, M., & Goshima, S. (2017). Life history traits and population dynamics of the invasive ascidian, *Ascidia aspersa*, on cultured scallops in Funka Bay, Hokkaido, northern Japan. *Journal of the Marine Biological Association of the United Kingdom*, 97(2), 387-399.
- Kim, Y. D., Lee, C., Shim, J. M., Kim, M., Kim, G. S., Choi, J.-S., An, W. G., & Nam, M.-M. (2014). A Study on the Growth of Juvenile *Patinopecten yessoensis* from Different Aquaculture Regions. *The Korean Journal of Malacology*, 30(4), 321-331.
- Kosaka, Y., & Ito, H. (2006). . "Chapter 22: Japan." *Developments in Aquaculture and Fisheries Science*.
- Kuriakose, S. (2017). Estimation of length weight relationship in fishes. *Course Manual Summer School on Advanced Methods for Fish Stock*

*Assessment and Fisheries Management. Lecture Note Series, L*, 215–220.

- Liu, F., Li, Y., Yu, H., Zhang, L., Hu, J., Bao, Z., & Wang, S. (2021). *MolluscDB: an integrated functional and evolutionary genomics database for the hyper-diverse animal phylum Mollusca*. 49(October 2020), 988–997.
- Matsushima, R., Uchida, H., Watanabe, R., Oikawa, H., Oogida, I., Kosaka, Y., Kanamori, M., Akamine, T., & Suzuki, T. (2018). Anatomical distribution of diarrhetic shellfish toxins (DSTs) in the Japanese scallop *Patinopecten yessoensis* and individual variability in scallops and *Mytilus edulis* mussels: Statistical considerations. *Toxins*, 10(10).
- Mau, A., Kangkan, A. L., & Ayubi, A. Al. (2023). Sebaran Panjang Kerang Kepah Di Desa Tanah Merah Kecamatan Kupang Tengah Kabupaten Kupang. Volume 4 (1). *E-ISSN: 2723-6536 E-ISSN: 2723-6536. April*, 145–153.
- Muchlisin, Z. A., Musman, M., & Siti Azizah, M. N. (2010). Length-weight relationships and condition factors of two threatened fishes, *Rasbora tawarensis* and *Poropuntius tawarensis*, endemic to Lake Laut Tawar, Aceh Province, Indonesia. *Journal of Applied Ichthyology*, 26(6), 949–953.
- Narejo, N. T., Rahmatullah, S. M., and Rashid, M. M. 2017. Length-weight relationship and relative condition factor (K<sub>n</sub>) of freshwater spiny eel, *Mastacembalus armatus* (Lacepede) from Mymensingh, Bangla-desh. *Indian Journal Fish*. 50(1): 81–87.
- Ogunola, O. S., Onada, O. A., Falaye, A. E., and Kunzman, A. 2017. Preliminary Investigation of Some Biological Aspects of Length-Weight Relationship and Condition Factor of Periwinkle (*Tympanotonus fuscatus*, Linnaeus 1758) from Okrika Estuary. *Global Journal of Science Frontier Research: Marine Science*. 17(1): 1-9 hal.
- Radiarta, I. N., & Saitoh, S. I. (2008). Satellite-derived measurements of spatial and temporal chlorophyll-a variability in Funka Bay, southwestern Hokkaido, Japan. *Estuarine, Coastal and Shelf Science*, 79(3), 400–408.
- Radiarta, I. N., Saitoh, S. I., & Miyazono, A. (2008). GIS-based multi-criteria evaluation models for identifying suitable sites for Japanese scallop (*Mizuhopecten yessoensis*) aquaculture in Funka Bay, southwestern Hokkaido, Japan. *Aquaculture*, 284(1–4), 127–135.
- Rochmady, R. (2012). Hubungan panjang bobot dan faktor kondisi kerang lumpur *Anodontia edentula*, Linnaeus 1758 di pulau Toba Kecamatan Napabalano, Kabupaten Muna. *Agrikan: Jurnal Agribisnis Perikanan*, 5(1), 1–8.

- Rochmady. 2012. Hubungan Panjang BObot dan Faktor Kondisi Kerang Lumpur *Anodontia Edentula* (Linnaeus, 1758) di Pulau Toba, Kecamatan Napabalano, Kabupaten Muna. *Jurnal Ilmiah Agribisnis dan Perikanan*. 5(1): 1-8 hal.
- Rohmah, A., & Muhsoni, F. F. (2020). Dinamika Populasi Kerang Tahu *Meretrix meretrix* Di Perairan Bangkalan Madura. *Juvenil: Jurnal Ilmiah Kelautan Dan Perikanan*, 1(3), 331-338.
- Sakurai, I., & Seto, M. (2000). Movement and orientation of the Japanese scallop *Patinopecten yessoensis* (Jay) in response to water flow. *Aquaculture*, 181(3-4), 269-279.
- Sharma, R., Venkateshvaran, K., and Purushothaman, C. S. 2005. Length-weight relationship and condition factor of *Perna viridis* (Linnaeus, 1758) and *Meretrix meretrix* (Linnaeus, 1758) from Mumbai waters. *Journal Indian Fish*. 32(2): 157 - 163.
- Silaban, R., Silubun, D. T., & Jamlean, A. A. R. (2021). Aspek Ekologi Dan Pertumbuhan Kerang Bulu (*Anadara antiquata*) Di Perairan Letman, KAabupaten Maluku Tenggara. *Jurnal Kelautan: Indonesian Journal of Marine Science and Technology*, 14(2), 120-131.
- Silina, A. V., & Zhukova, N. V. (2007). Growth variability and feeding of scallop *Patinopecten yessoensis* on different bottom sediments: Evidence from fatty acid analysis. *Journal of Experimental Marine Biology and Ecology*, 348(1-2), 46-59.
- Studi, P., Kelautan, I., Perikanan, F., & New, P. (2012). *Studi Bioekologi Kerang Simping ( Amusium pleuronectes ) di Perairan Semarang dan Kendal Diponegoro Kampus Tembalang , Semarang , 110-117.*
- Taufan, W.T., Anggoro, S., D., dan Widowati, I. 2016. Bioekologi Kerang Simping *Amusium pleuronectes* di Perairan Kabupaten Brebes. *Prosiding Seminar Nasional Tahunan Ke V Hasil-Hasil Penelitian Perikanan dan Kelautan*. 581-595.
- Uki, N. (2006). Stock enhancement of the Japanese scallop *Patinopecten yessoensis* in Hokkaido. *Fisheries Research*, 80(1), 62-66.
- Ventilla, R. F. (1982). The Scallop Industry in Japan. *Advances in Marine Biology*, 20(C), 309-382.
- Wanimbo, E., & Kalor, J. D. (2019). Morfometrik Kerang Polymesoda erosa di Perairan Teluk Youtefa Jayapura Papua. *ACROPORA: Jurnal Ilmu Kelautan Dan Perikanan Papua*, 1(2).
- Wiwiet Teguh Taufan, Anggoro, S., & Widowati, I. (2016). Bioekologi

Kerang Simping ( *Amusium pleuronectes* ) Di Perairan Kabupaten Brebes. *Prosiding Seminar Nasional Tahunan Ke V Hasil-Hasil Penelitian Perikanan Dan Kelautan, F1 07*, 581-595.

Woli, K. P., Nagumo, T., & Hatano, R. (2002). Evaluating impact of land use and N budgets on stream water quality in Hokkaido, Japan. *Nutrient Cycling in Agroecosystems*, 63(2-3), 175-184.

Yu, Z., Liu, C., Wang, F., Xue, Z., Zhang, A., Lu, G., Wang, L., & Song, L. (2019). Diversity and annual variation of phytoplankton community in Yesso scallop (*Patinopecten yessoensis*) farming waters of North Yellow Sea of China. *Aquaculture*, 511(June), 734266.

Zabarun, A., Bahtiar, & Haslianti. (2016). Hubungan panjang berat , faktor kondisi dan rasio berat daging Kerang Pasir ( *Modiolus modulaides* ) di perairan Bungkutoko Kota Kendari. *Jurnal Manajemen Sumberdaya Perairan*, 2(1), 21-32.

