

DAFTAR PUSTAKA

- Abida, I. W., Wahyuni, E. A., & Efendy, M. (2014). Hubungan Panjang Berat Lorjuk (*Solen spp*) di Perairan Pesisir Pantai Selatan Pulau Madura. *Jurnal Kelautan*, 7(1), 26–32. <http://journal.trunojoyo.ac.id/jurnalkelautan>
- Aya, F. A., Hidaka, Y., & Kudo, I. (2014). Clearance rates and ingestion efficiency of the Japanese scallop *Patinopecten yessoensis*. *Plankton and Benthos Research*, 8(3), 134–140. <https://doi.org/10.3800/pbr.8.134>
- Aya, F. A., & Kudo, I. (2007). Isotopic determination of Japanese scallop *Patinopecten* (*Mizuhopecten*) *yessoensis* (Jay) tissues shows habitat-related differences in food sources. *Journal of Shellfish Research*, 26(2), 295–302. [https://doi.org/10.2983/0730-8000\(2007\)26\[295:IDOJSP\]2.0.CO;2](https://doi.org/10.2983/0730-8000(2007)26[295:IDOJSP]2.0.CO;2)
- Bazikalova, A.Ya., Age and growth rate of *Pecten yessoensis* (Jay), *Izv. Akad. Nauk SSSR, Otd. Mat. Estestv. Nauk*, 1934, nos. 2–3, pp. 389–394.
- Buban, I. C. R., & Soliman, V. S. (2019). Review of scallop grow-out methods in tropical and temperate marine waters. *AACL Bioflux*, 12(2), 426–436.
- Coleman, S., Kiffney, T., Tanaka, K. R., Morse, D., & Brady, D. C. (2022). Meta-analysis of growth and mortality rates of net cultured sea scallops across the Northwest Atlantic. *Aquaculture*, 546(August 2021), 737392. <https://doi.org/10.1016/j.aquaculture.2021.737392>
- Dewi Pratiwi Eka Sari, F. (2019). Aspek Morfometri Kerang Darah (*Anadara Granosa L.*) Hasil Budidaya Di Perairan Desa Sukal, Kabupaten Bangka Barat. *Prosiding Seminar Hukum Dan Publikasi Nasional (Serumpun) I*, 978–623.
- Firdaus, Q. Y., & Safitri, N. M. (2023). Analisis Pertumbuhan Dan Kepadatan Kerang Hijau (*Perna Viridis*) Pada Tali Gantung Karamba Apung Kerang Hijau (*Kakh*) Di Laut Banyuurip. 6, 281–293.
- Grant, J., Emerson, C. W., Mallet, A., & Carver, C. (2003). Growth advantages of ear hanging compared to cage culture for sea scallops, *Placopecten magellanicus*. *Aquaculture*, 217(1–4), 301–323. [https://doi.org/10.1016/S0044-8486\(02\)00360-5](https://doi.org/10.1016/S0044-8486(02)00360-5)
- Hamsiah, Asmidar, Hasrun, & Kasmawati. (2018). Hubungan Panjang Berat dan Sebaran Ukuran Panjang Kerang Bakalang (*Marcia Hiantina Lamarck*) di Perairan Pesisir Labakkang, Kabupaten Pangkep Length weight relationship and size distribution long shells bakalang (*Marcia hiantina Lamarck*) in the coasta. *Torani: Journal of Fisheries and Marine Science*, 2(1), 23–31.
- Hartoyo Notonegoro, F. D. P. (2022). Morfometri Kerang *Anadara Granosa* Di Perairan Sukal Bangka Barat *Morfometric Of Anadara Granosa Shellfish In Sukal Waters, West*. 4(April), 18–22.
- Haryatik, R. . D., Hafiluddin, & Farid, A. (2013). Hubungan Panjang Berat dan Morfometrik Kerang Pisau (*Solen grandis*) di Perairan Prenduan Kabupaten Sumenep Madura. 3(1), 443–444.
- Herliantos, Bonte, S. B., Rosmianto, & Salim, G. (2012). Pengukuran Morfometri Kerang Kapah (*Meretrix lyrata*) Di Pantai Amal Lama Kota Tarakan. *Jurnal*

- Harpodon Borneo*, 5(2), 91–103.
- Indraswari, A. G. M., & Litaay, M. (2014). Morfometri Kerang Tahu Meretrix meretrix Linnaeus , 1758 Di Pasar Rakyat Makassar. *Berita Biologi*, 13(April), 137–142.
- Indraswari, A., Litaay, M., & Soekendarsi, E. (2014). Morfometri Kerang Tahu Meretrix meretrix Linnaeus, 1758 DI PASAR RAKYAT MAKASSAR [Morphometric of White Shell Meretrix meretrix Linnaeus, 1758 From Local Markets Makassar]. *Berita Biologi*, 13(2), 137–142.
- Kanamori, M., Baba, K., Natsuike, M., & Goshima, S. (2017). Life history traits and population dynamics of the invasive ascidian, *Ascidiella aspersa*, on cultured scallops in Funka Bay, Hokkaido, northern Japan. *Journal of the Marine Biological Association of the United Kingdom*, 97(2), 387–399. <https://doi.org/10.1017/S0025315416000497>
- Komala, R. D. (2017). Jurnal Fakultas Ilmu Terapan Universitas Telkom. *Jurnal Fakultas Ilmu Terapan Universitas Telkom*, 3(2), 330–337.
- Kosaka, Y. (2016). Scallop Fisheries and Aquaculture in Japan. In *Developments in Aquaculture and Fisheries Science* (Vol. 40). Elsevier B.V. <https://doi.org/10.1016/B978-0-444-62710-0.00021-3>
- Liu, Y., Saitoh, S. I., Igarashi, H., & Hirawake, T. (2014). The regional impacts of climate change on coastal environments and the aquaculture of Japanese scallops in northeast Asia: Case studies from Dalian, China, and Funka Bay, Japan. *International Journal of Remote Sensing*, 35(11–12), 4422–4440. <https://doi.org/10.1080/01431161.2014.916435>
- Lysenko, V. N., Zharikov, V. V., Lebedev, A. M., & Sokolenko, D. A. (2017). Distribution of the Yesso Scallop, *Mizuhopecten yessoensis* (Jay, 1857) (Bivalvia: Pectinidae) in the southern part of the Far Eastern Marine Reserve. *Russian Journal of Marine Biology*, 43(4), 302–311. <https://doi.org/10.1134/S1063074017040083>
- Mariani, S., Piccari, F., & De Matthaeis, E. (2002). Shell morphology in *Cerastoderma* spp. (Bivalvia: Cardiidae) and its significance for adaptation to tidal and non-tidal coastal habitats. *Journal of the Marine Biological Association of the United Kingdom*, 82(3), 483–490. <https://doi.org/10.1017/S0025315402005751>
- Matsushima, R., Uchida, H., Watanabe, R., Oikawa, H., Oogida, I., Kosaka, Y., Kanamori, M., Akamine, T., & Suzuki, T. (2018). Anatomical distribution of diarrhetic shellfish toxins (DSTs) in the Japanese scallop *Patinopecten yessoensis* and individual variability in scallops and *Mytilus edulis* mussels: Statistical considerations. *Toxins*, 10(10). <https://doi.org/10.3390/toxins10100395>
- Mulki, A. B. R., C. A. S. dan J. S. (2014). Variasi Ukuran Kerang Darah (*Anadara granosa*) di Perairan Pesisir Kecamatan Genuk Kota Semarang. *Journal of Marine Research*, 122–131. <http://ejournal-s1.undip.ac.id/index.php/jmr%0AVariasi>
- Mustamu, G., Lumingas, L. J., & Lohoo, A. V. (2014). Kepadatan, pola sebaran, dan morfometrik kerang kotak *Septifer bilocularis* (Linnaeus, 1758) pada rata-rata terumbu di Tanjung lampangi, Minahasa selatan. *Jurnal Ilmiah*

- Platax*, 2(1), 8–18.
- Nursalim, H. R., Suprijanto, J., & Widowati, I. (2012). Online di : <http://ejournal-s1.undip.ac.id/index.php/jmr> Studi Bioekologi Kerang Simpson (*Amusium pleuronectes*) di Perairan Semarang dan Kendal Diponegoro Kampus Tembalang , Semarang 50275 Telp / Fax . 024-7474698 Abstrak PENDAHULUAN Amusium pleuronecte. *Journal Of Marine Research*, 1(1), 110–117.
- Okayoshi, T. (2013). 岡吉 倫弘 1) , 三幡 輝久 1) , 渡辺 千聡 1) , 根尾 昌志 1). *Jurnal Rekayasa*, 6(1), 28–36.
- Pahri, S. D. R., Mohamed, A. F., & Samat, A. (2016). Preliminary water quality study in cockle farming area in Malaysia: A case study in Jeram, Selangor. *AAFL Bioflux*, 9(2), 316–325.
- Radiarta, I. N., Saitoh, S. I., & Miyazono, A. (2008). GIS-based multi-criteria evaluation models for identifying suitable sites for Japanese scallop (*Mizuhopecten yessoensis*) aquaculture in Funka Bay, southwestern Hokkaido, Japan. *Aquaculture*, 284(1–4), 127–135. <https://doi.org/10.1016/j.aquaculture.2008.07.048>
- Radiarta, I. N., Saitoh, S., & Yasui, H. (2011). *Aquaculture site selection for Japanese kelp (Laminaria japonica) in southern Hokkaido , Japan , using satellite remote sensing and GIS-based models*. 68, 773–780.
- Sangare, N., Lo-Yat, A., Moullac, G. Le, Pecquerie, L., Thomas, Y., Lefebvre, S., Gendre, R. Le, Beliaeff, B., & Andréfouët, S. (2020). Impact of environmental variability on *Pinctada margaritifera* life-history traits: A full life cycle deb modeling approach. *Ecological Modelling*, 423(February). <https://doi.org/10.1016/j.ecolmodel.2020.109006>
- Silaban, R., Dobo, J., & Rahanabun, G. (2022). Proporsi Morfometrik dan Pola Pertumbuhan Kerang Darah (*Anadara granosa*) di Daerah Intertidal, Kota Tual. *Jurnal Kelautan: Indonesian Journal of Marine Science and Technology*, 15(2), 143–152. <https://doi.org/10.21107/jk.v15i2.13759>
- Silina, A. V., & Zhukova, N. V. (2007). Growth variability and feeding of scallop *Patinopecten yessoensis* on different bottom sediments: Evidence from fatty acid analysis. *Journal of Experimental Marine Biology and Ecology*, 348(1–2), 46–59. <https://doi.org/10.1016/j.jembe.2007.03.018>
- Sumtaki, K., Kalesaran, O. J., & Lumenta, C. (2018). Telaah morfometrik *Pinctada margaritifera* untuk pengembangan usaha budidaya (Morphometric study of *Pinctada margaritifera* for the development of Pearl Aquaculture). *Budidaya Perairan Januari*, 6(1), 15–24.
- Torang J. P. Silalahi¹, Ockstan J. Kalesaran², Cyska Lumenta², Winda M. Mingkid², E. L. A. N. (2022). Karakteristik morfometrik kerang mutiara *Pinctada margaritifera* di Perairan Arakan, Mantehage Bango dan Talengen Provinsi Sulawesi Utara untuk kelayakan budidaya. 10(1), 10–20. <https://doi.org/10.14341/conf05-08.09.22-191>
- Torang J. P Silalahi, C. L. (2022). Morfometrik cangkang kerang mutiara *Pinctada margaritifera* untuk budidaya perairan. *Jurnal Moluska Indonesia*, 6(2), 54–58.
- Uki, N. (2006). Stock enhancement of the Japanese scallop *Patinopecten*

- yessoensis in Hokkaido. *Fisheries Research*, 80(1), 62–66.
<https://doi.org/10.1016/j.fishres.2006.03.013>
- Ulfa Apriliana, D., & Ambarwati, R. (2018). Karakter Morfologi dan Morfometri Kerang Eres dan Jubing (*Bivalvia: Pharidae*). *LenteraBio*, 7(3), 209–213.
<http://ejournal.unesa.ac.id/index.php/lenterabio>
- Uswatul Hasan, Hesti Wahyuningsih, dan E. J. (2014). Kepadatan dan Pola Pertumbuhan Kerang Lokan (*Geloina erosa*, Solander 1786) di Ekosistem Mangrove Belawan. *Jurnal Perikanan Dan Kelautan, Solander 1786*, 42–49.
- Wanimbo, E., & Kalor, J. D. (2019). Morfometrik Kerang Polymesoda erosa di Perairan Teluk Youtefa Jayapura Papua. *ACROPORA: Jurnal Ilmu Kelautan Dan Perikanan Papua*, 1(2). <https://doi.org/10.31957/acr.v1i2.930>
- Woli, K. P., Nagumo, T., & Hatano, R. (2002). Evaluating impact of land use and N budgets on stream water quality in Hokkaido, Japan. *Nutrient Cycling in Agroecosystems*, 63(2–3), 175–184.
<https://doi.org/10.1023/A:1021146707412>
- Zhao, L., Higuchi, T., Kanamori, M., Natsuike, M., Misaka, N., Murakami-Sugihara, N., Tanaka, K., & Shirai, K. (2021). Identification of timing of scallop morphological deformity and mortality from shell oxygen isotope records. *Marine Environmental Research*, 163, 105149.
<https://doi.org/10.1016/j.marenvres.2020.105149>

