

## DAFTAR PUSTAKA

- Agassiz, A. 1864. Synopsis of the echinoids collected by Dr. W. Stimpson on the North Pacific Exploring Expedition under the command of Captains Ringgold and Rodgers. *Proceedings of the Academy of Natural Sciences of Philadelphia*. 15 (1863), 352-361.
- Agatsuma, Y. 2013. *Strongylocentrotus intermedius*. In Sea Urchins: Biology and Ecology., Elsevier. pp. 438-447.
- Agatsuma, Y. 2014. Population dynamics of edible sea urchins associated with variability of seaweed beds in northern Japan. Terrapub.
- Aguera, A., Trommelen, M., Burrows, F., Jansen, J. M., Schellekens, T., & Smaal, A. 2012. Winter feeding activity of the common starfish (*Asterias rubens* L.): The role of temperature and shading. *Journal of Sea Research*, 72, 106-112.
- Andrews, et al. 2010. *Japanese Scallop Hanging and Seabed Enhanced Fisheries*. Hokkaido, Jepang.
- Anggorowati. 2008. *Biologi Laut, Suatu Pendekatan Ekologis*. Jakarta: PT. Gramedia.
- Asai, Naoki. 2000. Sex pheromones of the hair crab *Erimacrus isenbeckii*. Part 1: Isolation and structures of novel ceramides. *Tetrahedron*. 56 (51): 9895-9899.
- Azuma, N., Kunihiro, Y., Sasaki, J., Mihara, E., Mihara, Y., Yasunaga, T., Jin, D., Abe, S. 2008. Genetic variation and population structure of hair crab (*Erimacrus isenbeckii*) in Japan inferred from mitochondrial DNA sequence analysis. *Marine Biotechnology*. 10 (1): 39-48.
- Borsa, P., Rolland, V., Daguin-Thiebaut, C. 2012. Genetics and taxonomy of Chilean smooth-shelled mussels, *Mytilus* spp. (Bivalvia: Mytilidae). *Comptes Rendus Biologies*. 335 (1): 51-61.
- Byrne, M., Morrice, M. G., & Wolf, B. 1997. Introduction of the northern Pacific asteroid *Asterias amurensis* to Tasmania: reproduction and current distribution. *Marine Biology*, 127, 673-685.
- Byrne, M., Andrew, N.L., Worthington, D. G., Brett, P. A. 1998. Reproduction in the diadematoid sea urchin *Centrostephanus rodgersii* in contrasting habitats along the coast of New South Wales, Australia. *Mar. Biol.* 132: 305-318
- Byrne, M., O Hara, T., & Lawrence, J. 2013. *Asterias amurensis*. In Lawrence, (Ed.), *Starfish: Biology and ecology of the asteroidea* (pp. 174-180). Baltimore, MD: JHU Press.
- Caro, A.U., & Castilla, J.C. 2004. Predator-inducible defenses and local intrapopulation variability of the intertidal mussel *Semimytilus algosus* in central Chile, *Mar. Ecol. Prog. Ser.* 276: 115-123.
- Chaliyenko, M. O. 2018. Features of group growth of the gray sea urchin (*Strongylocentrotus intermedius*) off the northwest coast of the Sea of Japan. *Izv. TINRO*, 194, 3-17.
- Chaliyenko, M. O., Kulepanov, V. N., & Matveev, V. I. 2018. The influence of some environmental factors on the growth of the gray sea urchin

- (*Strongylocentrotus intermedius*) off the northwest coast of the Sea of Japan. *Izv. TINRO*, 195, 111–127.
- Chaliyenko, M. O., Kalinina, M. V., Kulepanov, V. N., Matveev, V. I. 2021. Size and Age at Maturity of the Sea Urchin *Strongylocentrotus intermedius* along the Northwestern Coast of the Sea of Japan. *Oceanology*, 61, 69–79.
- Chuchukalo, V. I., Nadtochy, V. A., Koblikov, V. N., Borilko, O. Y. 2011. Diet and some ecological features of the most widespread commercial crab species in the northwestern Sea of Japan in early spring. *Russian Journal of Marine Biology*. 37 (7): 558–569.
- Cinar, M. E. 2006. Serpulid species (Polychaeta: Serpulidae) from the Levantine coast of Turkey (eastern Mediterranean), with special emphasis on alien species. *Aquatic Invasions*. 1(4): 223–240
- Cunningham, C. W., Blackstone, N. W., & Buss, L. W. 1992. Evolution of king crab from hermit crab ancestors. *Nature*, 355: 539–542.
- Dautov, S., Dautova, T., & Kashenko, S. 2020. Towards a scientific-based farming of sea urchins: First steps in the cultivation of *Diadema setosum*, *Diadema savignyi* and *Mesocentrotus nudus*. *APN Science Bulletin*.
- Davydov, P. V.; Shubrayvi, O. I.; Vassetzky, S. G. 1990. The starfish *Asterina pectinifera* (Müller et Troschel, 1842). *Animal Species for Developmental Studies*. pp. 287–311.
- Evdokimov, V. V., & Matrosova, I. V. 2012. Reproductive biology of sea urchins *Strongylocentrotus intermedius* and *Strongylocentrotus nudus*. *Pac. Med. J.* 2, 105–110.
- FAO (Food and Agriculture Organization). 2006. *Fisheries global information systems*. FAO, Rome.
- FAO (Food and Agriculture Organization). 2009. *The State of World Fisheries and Aquaculture, 2008*, pp. 196. FAO, Rome.
- Gabaev, D. D. 2018. Ecology of the Starfish *Asterias amurensis* (Lutken, 1871) in Russia's East. *American Journal of Marine Science*, 6(1), 1–19.
- Gardner, J. P. A., & Thompson, R. J. The effects of coastal and estuarine conditions on the physiology and survivorship of the mussels *Mytilus edulis*, *M. trossulus* and their hybrids. *J Exp Mar Bio Ecol.* 265 (2): 119–140.
- Gillespie, Graham E., Bower, Susan M., Marcus, Kerry L., Kieser, Dorothee. 2012. Biological synopses for three exotic molluscs, Manila Clam (*Venerupis philippinarum*), Pacific Oyster (*Crassostrea gigas*) and Japanese Scallop (*Mizuhopecten yessoensis*) licensed for Aquaculture in British Columbia. *Canadian Science Advisory Secretariat, Fisheries and Oceans Canada, Ottawa, Ontario*. Pp. 1–97.
- Guillou, M., Joly-Turquin, G., Leyzour, S., et al., 2012. Factors controlling juvenile growth and population structure of the *Asterias rubens* in intertidal habitats: field and experimental approaches, *J. Mar. Biol. Ass. U.K.* 92: 367–378.
- Hayward, P. J., & Ryland, J. S. 1995. *Handbook of the marine fauna of north-west Europe*. Oxford: Oxford University Press.

- Hansen, T. 2002. *Temperature preferences of the Red king crab (Paralithodes camtschaticus)*. University of Tromsø, Norway: Norwegian College of Fishery Science, 86 pp.
- Haswell, W. A. 1883. On some new Australian tubicolous annelida. *Proceedings of the Linnean Society of New South Wales*, 7: 633–638.
- Hemmingsen, W., Jansen, P. A., & MacKenzie, K. 2005. Crabs, leeches and trypanosomes: an unholy trinity. *Marine Pollution Bulletin*, 50 (3): 336–339.
- Hood, D. W., Fisher, V., Nebert, D., Feder, H., Mueller, G. L., Burrell, D., Boisseau, D., Goering, J. J., Sharma, G. D., Kresge, D. T., Fison, S. R. 1974. *Environmental study of the marine environment near Nome, Alaska*. University of Alaska, Institute of Marine Science, 265 pp.
- Hutson, K. S., Ross, D. J., Day, R. W., & Ahern, J. J. 2005. Australian scallops do not recognise the introduced predatory seastar *Asterias amurensis*. *Marine Ecology Progress Series*, 298, 305–309.
- Hwang, *et al.* 2015. Invertebrates fauna in the intertidal regions of Yubudo Island, South Korea. *Journal of Asia-Pacific Biodiversity*, 8(1): 66–71.
- Ito, H. 1991. Japan. In: Shumway S (ed) *Scallop: biology, ecology and aquaculture. Development in aquaculture and fisheries science*, vol 21. Elsevier, Amsterdam, pp 1017–1055.
- Japan Fisheries Association. 2020. *Buku Teks Untuk Tes Keterampilan Perikanan (Budidaya)*. Hokkaido, Jepang.
- Jay, J. C. 1857. Report on the shells collected by the Japan Expedition together with a list of Japan shells. In: Perry, M. C.; Hawks, F. L. (eds). *Narrative of the expedition of an American squadron to the China Sea and Japan, performed in the year 1852, 1853 and 1854, under the command of Commodore M. C. Perry, United States Navy, by order of the Government of the United States*. Vol. 2. Tucker, Washington, pp. 289–297, pls. 1–5.
- Jewett, S. C., & Onuf, C. P. 1988. Habitat suitability index models: Red king crab. U.S. Fish & Wildl. Ser. Biol. Rep., National Wetlands Research Center, Slidell. Louisiana. 34 pp.
- Kado, R., Suzuki, J., Nanba, N., & Ogawa, H. 2009. Reproduction, growth, and feeding habits of *Balanus rostratus* Hoek transplanted into Okkirai Bay in northern Honshu, Japan, and prospective problems for its aquaculture. *Sessile Organisms*, 26, 1–10.
- Kado, R., Suzuki, J., Suzuki, Y., Nanba, N., & Ogawa, H. 2009. Reproductive cycle, larval distribution, settlement and initial growth of the barnacle *Balanus rostratus* in Mutsu Bay, northern Japan. *Nippon Suisan Gakkaishi*, 75, 432–442.
- Kai, Y., Xiaoqi, Z., Hui, L., Xiumei, Z., Qun, L. 2003. Selectivity and tolerance of sea urchin (*Hemicentrotus pulcherrimus*) to environmental change. *Chin. J. App. Ecol.* 14: 409– 412.
- Kashenko, S. D. 2003. The Reaction of the Starfish *Asterias amurensis* and *Patiria pectinifera* (Asteroidea) from Vostok Bay (Sea of Japan) to a Salinity Decrease. *Russian Journal of Marine Biology*, 29(2): 110–114.

- Kashenko, S. D. 2005. Chronology of Development in the Starfish *Asterina pectinifera* from Vostok Bay, Sea of Japan. *Russian Journal of Marine Biology*, 31(4): 261-264.
- Kashenko, S. D. 2006. The combined effect of temperature and salinity on development of the sea star *Asterina pectinifera*. *Russian Journal of Marine Biology*, 32(1): 37-44.
- Kashenko, S. D. 2012. Resistance of sea urchins *Strongylocentrotus nudus* and *S. intermedius* to experimental changes in the salinity of sea water. *Russ. J. Mar. Biol.* 38, 395–399.
- Kasyanov, V. L., Medvedeva, L. A., Yakovlev, U. M., & Yakovlev, S. N. 1980. *Breeding of Echinoderms and Marine Clams*. Nauka, Moscow.
- Katolikova, M., Khaitov, V., Vainola, R., Gantsevich, M., & Strelkov, P. 2016. Genetic, ecological and morphological distinctness of the blue mussels *Mytilus trossulus* Gould and *M. edulis* L. in the White Sea. *PLoS One*, 11(4), e0152963.
- Kim, Sang Gu. 2011. Hemocyanin-derived phenoloxidase activity with broad temperature stability extending into the cold environment in hemocytes of the hair crab *Erimacrus isenbeckii*. *Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology*. 159 (2): 103–108.
- Klibansky, L. K. J., & McCartney, M. A. 2014. Conspecific sperm precedence is a reproductive barrier between free spawning marine mussels in the northwest Atlantic *Mytilus* hybrid zone. *PLoS One*. 9(9).
- Kott, P. A. 1985. The Australian Ascidiacea. Part 1. Phlebobranchia and Stolidobranchia. *Memoirs of the Queensland Museum*, 23, 1–440.
- Lee, C., & Ko, H. S. 2010. Early zoeal stages of edible crab *Erimacrus isenbeckii* (Brandt, 1848) (Crustacea: Decapoda: Brachyura: Cheiragonidae) and a comparison with other cheiragonid zoeae. *Animal cells and systems*, 14(4), 323-331.
- Lynch, S. A., Darmody, G., O'Dwyer, K., Gallagher, M. C., Nolan, S., McAllen, R., & Culloty, S. C. 2016. Biology of the invasive ascidian *Asciidiella aspersa* in its native habitat: Reproductive patterns and parasite load. *Estuarine, Coastal and Shelf Science*, 181, 249-255.
- Lutken, C. 1871. Fortsatte kritiske og beskrivende Bidrag til Kundskab om Söstjerneerne (Asteriderne). *Videnskabelige Meddelelser fra Dansk Naturhistorisk Forening*, 23, 227-304.
- Manchenko, G. P., & Serov, O. L. 1976. Electrophoretic spectra of isomerases, transferases and oxidoreductases of the starfish *Patina pectinifera*. *Fisheries and Marine Service*.
- Mastrototaro, F., D'Onghia, G., & Tursi, A. 2008. Spatial and seasonal distribution of ascidians in a semi-enclosed basin of the Mediterranean Sea. *Journal of the Marine Biological Association of the United Kingdom*, 8: 1053–61.
- Mathiesen, S. S., Thyrring, J., Hemmer-Hansen, J., Berge, J., Sukhotin, A., Leopold, P., Bekaert, M., Sejrs, M. K., Nielsen, E. 2016. Genetic diversity and connectivity within *Mytilus* spp. in the subarctic and Arctic. *Evolutionary Applications*. 10 (1): 39–55.

- Minchin, Dan. 2003. Introductions: some biological and ecological characteristics of scallops. *Aquatic Living Resources* 16: 521-532.
- Miranda, M. B. B., Innes, D. J., & Thompson, R. J. 2010. Incomplete Reproductive isolation in the blue mussel (*Mytilus edulis* and *M. trossulus*) hybrid zone in the northwest Atlantic: Role of gamete interactions and larval viability. *Biol Bull.* 218(3): 266-281.
- Miyahara, D., & Kado, R. 2022. Growth, feeding, and larval density of *Balanus rostratus* in Ofunato Bay, northern Honshu, Japan, compared with those in adjacent Mutsu Bay, Okirai Bay and Shizugawa Bay. *Sessile Organisms*, 39(1), 1-13.
- Moreau, V., Tremblay, R., Bourget, E. 2005. Distribution of *Mytilus edulis* and *M. trossulus* on the Gaspé coast in relation to spatial scale. *J Shellfish Res.* 24(2): 545-551.
- Morton, B., & Dinesen, G. E. 2011. The biology and functional morphology of *Modiolarca subpicta* (Bivalvia: Mytilidae: Musculinae), epizoically symbiotic with *Asciidiella aspersa* (Urochordata: Ascidiacea), from the Kattegat, northern Jutland, Denmark. *Journal of the Marine Biological Association of the United Kingdom*, 91(8), 1637-1649.
- Murabe, N., H. Hatoyama, K. Mieko, H. Kaneko, Y. Nakajima. 2007. Adhesive papillae on the brachiolar arms of brachiolaria larvae in two starfishes, *Asterina pectinifera* and *Asterias amurensis*, are sensors for metamorphic inducing factors(s). *Development, Growth, and Differentiation*, 49(8): 647-656.
- Nagao, Jiro. 1999. Embryonic development of the hair crab *Erimacrus Isenbeckii*. *Journal of Crustacean Biology*. 19 (1): 77-83.
- Nedved, B. T., & Hadfield, M. G. 2008. *Hydroides elegans* (Annelida: Polychaeta): a model for biofouling research.
- Nydam, M. L., Nichols, C. L., & Lambert, G. 2022. First record of the ascidian *Asciidiella aspersa* (Müller, 1776) in southern California. *BioInvasions Record*, 11(2).
- Paik, S., H. Park, & S. Yi, S. Yun. 2005. Developmental duration and morphology of the sea star *Asterias amurensis* in Tongyeong, Korea. *Ocean Science Journal*, 40(3): 65-70.
- Pearse, J. S., & Cameron, R. A. 1991. Echinodermata: Echinoidea. In: Giese AC, Pearse JS, Pearse VB (eds). *Reproduction of Marine Invertebrates. Vol. 6.* The Boxwood Press, Pacific Grove, California. 514-662.
- Powell, G. C., & Nickerson, R. B. 1965. Reproduction of king crabs *Paralithodes camtschatica* (Tilesius). *J. Fish. Res. Bd. Can*, 22 (1): 101-111.
- Qiu, J.W., & Qian, P. 1997. Combined effects of salinity, temperature and food on early development of the polychaete *Hydroides elegans*. *Marine Ecology Progress Series*. 152: 79-88
- Qiu, J. W., Tremblay, R., Bourget, E. 2002. Ontogenetic changes in hyposaline tolerance in the mussels *Mytilus edulis* and *M. trossulus*: Implications for distribution. *Mar Ecol Prog Ser.* 228: 143-152.
- Radiarta, I. N., & Saitoh, S. I. 2009. Biophysical models for Japanese scallop, *Mizuhopecten yessoensis*, aquaculture site selection in Funka Bay,

- Hokkaido, Japan, using remotely sensed data and geographic information system. *Aquaculture international*, 17(5), 403-419.
- Radiarta, I. N., Saitoh, S. I., & Miyazono, A. (2008). GIS-based multi-criteria evaluation models for identifying suitable sites for Japanese scallop (*Mizuhopecten yessoensis*) aquaculture in Funka Bay, southwestern Hokkaido, Japan. *Aquaculture*, 284(1-4), 127-135.
- Ristante, A., Yanti, A. H., & Setyawati, T. R. 2018. Sea Urchin (Echinoidea) Distribution and Abundance in the Intertidal Zone of Bengkayang Regency. *Biosaintifika: Journal of Biology & Biology Education*, 10(1), 32-40.
- Rolando Bastida-Zavala, J., & ten Hove, H. 2002. Revision of Hydroides gunnerus, 1768 (Polychaeta: Serpulidae) from the western Atlantic region. *Beaufortia*, 52(9), 103-178.
- Ruppert, E. E., Fox, R. S., & Barnes, R. D. 2004. Invertebrate Zoology. A functional evolutionary approach. 7th Ed. Brooks/Cole, Thomson Learning learning, Inc. 990 p.
- Rusanowski, P. C., Gardner, L. A., & Jewett, S. C. 1987. Nome offshore placer project. Annual report. Prepared for Inspiration Gold, Inc. Nome, Alaska. 212 pp.
- Sasaki, J. 1993. Early larval stages of the hair crab *Erimacrus isenbeckii* (Brandt) (Brachyura: Atelecyclidae), with special reference to its hatching process. *Journal of Crustacean Biology*. 13 (3): 511-522.
- Sato, M. *et al.* 2005. Development of microsatellite markers for Japanese scallop (*Mizuhopecten yessoensis*) and their application to a population 22 genetic study, *Marine Biotechnology*, 7(6), pp. 713-728.
- Shumway, S. E. 1991. Scallops: biology, ecology and aquaculture. Development in aquaculture and fisheries science. *Elsevier*. Amsterdam. Vol 21.
- Silina, A. V. 2018. Sex change in scallop *Patinopecten yessoensis*: response to population composition?. *PeerJ*, 6.
- Silina, A. V., & Ovsyannikova, I. I. 1998. Barnacle *Balanus rostratus* in the Northwestern Part of the Sea of Japan. *Ophelia*, 4, pp. 28-35.
- Silina, A. V., & Ovsyannikova, I. I. 1999. Growth of the barnacle *Balanus rostratus* in the Sea of Japan. *Russian Journal of Marine Biology*, 25(1), 18-22.
- Silina, A. V., & Ovsyannikova, I. I. 2000. Variability in morphology of the shell of the barnacle, *Balanus rostratus*, under different conditions of growth (Cirripedia, Thoracica). *Crustaceana*, 519-524.
- Stevens, B. G. 2014. *King Crabs of the World: Biology and Fisheries Management*. CRC Press. pp. 3-7.
- Sundet, J. H., Kuzmin, S. A., Hjelset, A. M., Nilsen, E. M. 2000. *Migration and migration patterns of red king crab (Paralithodes camtschaticus) in the southern Barents Sea, Varanger Area Crab*. In: 19th Lowell Wakefield Fisheries Symposium: Crabs in Cold Water Regions: Biology, Management, and Economics. Anchorage, Alaska, USA.
- Suryanti, S., Fatimah, P. N. P. N., & Rudiyananti, S. 2020. Morfologi, Anatomi dan Indeks Ekologi Bulu Babi di Pantai Sepanjang, Kabupaten Gunungkidul, Yogyakarta. *Buletin Oseanografi Marina*, 9(2), 93-103.

- Takagi, S., Murata, Y., Inomata, E., Endo, H., Aoki, M. N., & Agatsuma, Y. 2018. Dietary Effect of Kelp (*Saccharina japonica*) on Gonad Quantity and Quality in Sea Urchins (*Mesocentrotus nudus*) Collected from a Barren Before the Fishing Season. *Journal of Shellfish Research*, 37(3), 659-669.
- Tarakanadha, B., Morrell, J. J., & Rao, K. S. 2004. Environmental impacts of preservative-treated wood. *Florida Center for Environmental Solutions, Orlando*, 320-335.
- Tavares, M., & Cleva, R. 2010. Trichopeltariidae (Crustacea, Decapoda, Brachyura), a new family and superfamily of eubrachyuran crabs with description of one new genus and five new species. *Papeis Avulsos Zool.* 50: 97-157.
- Teshirogi, W. and Ishida, S. 1978. Early development of the starfish *Asterina pectinifera*, especially on the cleavage patterns, blastomere fusion, wrinkled morula, and wrinkled blastula. *Rep. Fukaura Mar. Bio. Lab*, 7, 2-24.
- Teshirogi, W., Ishida, S., and Kodama, M. 1980. Early development of the starfish eggs (*Asterina pectinifera*) inseminated with various sperm concentrations in oocyte maturation process. *Rep. Fukaura Mar. Bio. Lab.*, 8, 2-13.
- Toro, J. E., Thompson, R. J., Innes, D. J. 2002. Reproductive isolation and reproductive output in two sympatric mussel species (*Mytilus edulis*, *M. trossulus*) and their hybrids from Newfoundland. *Mar Biol.* 141 (5): 897-909.
- Wang, J., & Huang, Z. 1993. Fouling polychaetes of Hong Kong and adjacent waters. *Asian Marine Biology*, 10: 1-12.
- Won, N. I., Kawamura, T., Takami, H., Watanabe, Y. 2013. Trophic structure in natural habitats of the abalone *Haliotis discus hannai* with distinct algal vegetation of kelp and crustose coralline algae: implication of ontogenetic niche shifts. *Fisheries Science*, 79(1): 87-97.
- Yamaguchi, A., Furumitsu, K., Yagishita, N., Kume, G. 2010. *Biology of herbivorous fish in the coastal areas of western Japan*. In: Ishimatsu A, Lie HJ (eds). *Coastal Environmental and Ecosystem Issues of the East China Sea*. TERRAPUB and Nagasaki University, Tokyo. 181-190.
- Yusup, D. S. 2009. Response of north pacific seaster *Asterias amurensis* to food attractant (Betaine), *Journal Biologi*. XVI (2): 52-56.
- Zhadan, P. M., & Vaschenko, M. A. 2019. Long-term study of behaviors of two cohabiting sea urchin species, *Mesocentrotus nudus* and *Strongylocentrotus intermedius*, under conditions of high food quantity and predation risk in situ. *PeerJ*, 7, e8087.
- Zolotarev, V. N. 2002. *Morphological differences in mussels of Mytilus edulis group*. Vestn Zhitomirskogo Derzhavnogo Univ Imi Franka. pp. 5-8.