

## DAFTAR PUSTAKA

- Admasu, T.D., Rae, M. and Stolzing, A. (2021) ‘Dissecting primary and secondary senescence to enable new senotherapeutic strategies.’, *Ageing research reviews*, 70, p. 101412.
- Alexandrov, N. V, Eelderink, C., Singh-Povel, C.M., Navis, G.J., et al. (2018) ‘Dietary protein sources and muscle mass over the life course: the lifelines cohort study’, *Nutrients*, 10(10).
- Anderson, L.J., Liu, H. and Garcia, J.M. (2017) ‘Sex differences in muscle wasting.’, *Advances in experimental medicine and biology*, 1043, pp. 153–197. Available at: [https://doi.org/10.1007/978-3-319-70178-3\\_9](https://doi.org/10.1007/978-3-319-70178-3_9).
- Aragon, A.A., Tipton, K.D. and Schoenfeld, B.J. (2023) ‘Age-related muscle anabolic resistance: inevitable or preventable?’, *Nutrition reviews*, 81(4)
- Asgari, M.R., Asghari, F., Ghods, A.A., Ghorbani, R., et al. (2017) ‘Incidence and severity of nausea and vomiting in a group of maintenance hemodialysis patients.’, *Journal of renal injury prevention*, 6(1), pp. 49–55.
- Beberashvili, I., Azar, A., Khatib, A., Abu Hamad, R., et al. (2023) ‘Sarcopenic obesity versus nonobese sarcopenia in hemodialysis patients: Differences in nutritional status, quality of life, and clinical outcomes.’, *Journal of renal nutrition: the official journal of the Council on Renal Nutrition of the National Kidney Foundation*, 33(1), pp. 147–156.
- Bernier-Jean, A., Prince, R.L., Lewis, J.R., Craig, J.C., et al. (2021) ‘Dietary plant and animal protein intake and decline in estimated glomerular filtration rate among elderly women: a 10-year longitudinal cohort study’, *Nephrology Dialysis Transplantation*, 36(9), pp. 1640–1647.
- Betiru, E.A., Mamo, E., Jara Boneya, D., Adem, A., et al. (2023) ‘Survival analysis and its predictors among hemodialysis patients at Saint Paul Hospital Millennium Medical College and Myungsung Christian Medical Center in Addis Ababa, Ethiopia, 2021.’, *International journal of nephrology and renovascular disease*, 16, pp. 59–71.
- Buckinx, F., Landi, F., Cesari, M., Fielding, R.A., et al. (2018) ‘Pitfalls in the measurement of muscle mass: a need for a reference standard’, *Journal of cachexia, sarcopenia and muscle*, 9(2), pp. 269–278.

- Campbell, K.L. and Carrero, J.J. (2016) 'Diet for the Management of Patients With Chronic Kidney Disease; It Is Not the Quantity, but the Quality That Matters.', *Journal of renal nutrition: the official journal of the Council on Renal Nutrition of the National Kidney Foundation*. United States, pp. 279–281. Available at: <https://doi.org/10.1053/j.jrn.2016.07.004>.
- Carbone, J.W. and Pasiakos, S.M. (2019) 'Dietary Protein and Muscle Mass: Translating Science to Application and Health Benefit.', *Nutrients*, 11(5).
- Cobo, G., Hecking, M., Port, F.K., Exner, I., et al. (2016) 'Sex and gender differences in chronic kidney disease: progression to end-stage renal disease and haemodialysis.', *Clinical science (London, England : 1979)*, 130(14), pp. 1147–1163. Available at: <https://doi.org/10.1042/CS20160047>.
- Crews, D.C., Kuczmarski, M.F., Miller III, E.R., Zonderman, A.B., et al. (2015) 'Dietary habits, poverty, and chronic kidney disease in an urban population', *Journal of Renal Nutrition*, 25(2), pp. 103–110.
- Cupisti, A., Brunori, G., Di Iorio, B.R., D'Alessandro, C., et al. (2018) 'Nutritional treatment of advanced CKD: twenty consensus statements.', *Journal of nephrology*, 31(4), pp. 457–473.
- Curtis, E., Litwic, A., Cooper, C. and Dennison, E. (2015) 'Determinants of muscle and bone aging.', *Journal of cellular physiology*, 230(11), pp. 2618–2625. Available at: <https://doi.org/10.1002/jcp.25001>.
- Damayanti, D. (2016) *Ilmu Gizi: Teori dan Aplikasi*. Edited by Hardinsyah and I.D.N. Supariasa. Jakarta: EGC.
- Deger, S.M., Hung, A.M., Gamboa, J.L., Siew, E.D., et al. (2017) 'Systemic inflammation is associated with exaggerated skeletal muscle protein catabolism in maintenance hemodialysis patients.', *JCI insight*, 2(22). Available at: <https://doi.org/10.1172/jci.insight.95185>.
- Denic, A., Lieske, J.C., Chakkera, H.A., Poggio, E.D., et al. (2017) 'The substantial loss of nephrons in healthy human kidneys with aging.', *Journal of the American Society of Nephrology : JASN*, 28(1), pp. 313–320.
- Dev, R., Hui, D., Chisholm, G., Delgado-Guay, M., et al. (2015) 'Hypermetabolism and symptom burden in advanced cancer patients evaluated in a cachexia clinic.', *Journal of cachexia, sarcopenia and muscle*, 6(1), pp. 95–98.

- Available at: <https://doi.org/10.1002/jcsm.12014>.
- Donini, L.M., Busetto, L., Bischoff, S.C., Cederholm, T., et al. (2022) 'Definition and diagnostic criteria for Sarcopenic Obesity: ESPEN and EASO Consensus Statement.', *Obesity facts*, 15(3), pp. 321–335.
- El-Kateb, S. and Davenport, A. (2016) 'Changes in intracellular water following hemodialysis treatment lead to changes in estimates of lean tissue using Bioimpedance Spectroscopy.', *Nutrition in clinical practice: official publication of the American Society for Parenteral and Enteral Nutrition*, 31(3), pp. 375–377.
- Fadili, W., Lahlou, L. and Laouad, I. (2023) 'Smoking in chronic hemodialysis patients: prevalence and cardiovascular morbidity', *Nephrologie & Thérapeutique*, 19(7), pp. 1–10.
- FAO (2011) 'Dietary protein quality evaluation in human nutrition', *FAO Food Nutr. Pap*, 92, pp. 1–66.
- Francaux, M. and Deldicque, L. (2019) 'Exercise and the control of muscle mass in human.', *Pflugers Archiv : European journal of physiology*, 471(3), pp. 397–411. Available at: <https://doi.org/10.1007/s00424-018-2217-x>.
- Frontera, W.R. and Ochala, J. (2015) 'Skeletal muscle: a brief review of structure and function.', *Calcified tissue international*, 96(3), pp. 183–195. Available at: <https://doi.org/10.1007/s00223-014-9915-y>.
- Garibotto, G., Sofia, A., Parodi, E.L., Ansaldi, F., et al. (2018) 'Effects of Low-Protein, and Supplemented Very Low-Protein Diets, on Muscle Protein Turnover in Patients With CKD', *Kidney International Reports*, 3(3), pp. 701–710.
- Goldberg, I. and Krause, I. (2016) 'The role of gender in chronic kidney disease', *Emj*, 1(2), pp. 58–64.
- Gollie, J.M., Harris-Love, M.O., Patel, S.S. and Argani, S. (2018) 'Chronic kidney disease: considerations for monitoring skeletal muscle health and prescribing resistance exercise.', *Clinical kidney journal*, 11(6), pp. 822–831.
- Guzman-Ventura, W. and Caballero-Alvarado, J. (2022) 'Survival of patients on chronic hemodialysis versus chronic peritoneal dialysis.', *Revista peruana de medicina experimental y salud publica*, 39(2), pp. 161–169.

- Haizlip, K.M., Harrison, B.C. and Leinwand, L.A. (2015) 'Sex-based differences in skeletal muscle kinetics and fiber-type composition', *Physiology*, 30(1), pp. 30–39.
- Hajj-Boutros, G., Karelis, A.D., Cefis, M., Morais, J.A., et al. (2022) 'Potential mechanisms involved in regulating muscle protein turnover after acute exercise: A brief review.', *Frontiers in physiology*, 13, p. 1106425.
- Hall, J.E. (2021) *Guyton and Hall Textbook of Medical Physiology*. 14th edn. Philadelphia: Elsevier.
- Hammer, G.D. and McPhee, S.J. (2019) *in Pathophysiology of Disease: An Introduction to Clinical Medicine*. 8th edn. New York, NY: McGraw-Hill Education.
- Hoppe, K., Schwermer, K., Kawka, A., Klysz, P., et al. (2018) 'Dialysis vintage stratified comparison of body composition, hydration and nutritional state in peritoneal dialysis and hemodialysis patients.', *Archives of medical science : AMS*, 14(4), pp. 807–817.
- Hsu, C.-C., Sun, C.-Y., Tsai, C.-Y., Chen, M.-Y., et al. (2021) 'Metabolism of Proteins and Amino Acids in Critical Illness: From Physiological Alterations to Relevant Clinical Practice.', *Journal of multidisciplinary healthcare*, 14, pp. 1107–1117. Available at: <https://doi.org/10.2147/JMDH.S306350>.
- Huh, Y. and Son, K.Y. (2022) 'Association between total protein intake and low muscle mass in Korean adults.', *BMC geriatrics*, 22(1), p. 319.
- Ikizler, T.A. (2013) 'A patient with CKD and poor nutritional status.', *Clinical journal of the American Society of Nephrology : CJASN*, 8(12), pp. 2174–2182. Available at: <https://doi.org/10.2215/CJN.04630513>.
- Ikizler, T.A., Burrowes, J.D., Byham-Gray, L.D., Campbell, K.L., et al. (2020) 'KDOQI Clinical Practice Guideline for Nutrition in CKD: 2020 Update.', *American journal of kidney diseases : the official journal of the National Kidney Foundation*, 76(3 Suppl 1), pp. S1–S107.
- Irwan (2016) *Epidemiologi Penyakit Tidak Menular*. Yogyakarta: Deepublish.
- Jain, R.K. and Vokes, T. (2017) 'Dual-energy X-ray absorptiometry', *Journal of Clinical Densitometry*, 20(3), pp. 291–303.
- Jamal, A., Babazono, A., Li, Y., Yoshida, S., et al. (2021) 'Quantifying regional

- and health care variations to identify ways to improve hemodialysis service quality and survival outcomes', *American Journal of Medical Quality*, 36(5), pp. 345–354.
- Jeznach-Steinhagen, A., Boniecka, I., Rymarz, A., Staszków, M., et al. (2023) 'Higher muscle mass and higher serum prealbumin levels are associated with better survival in hemodialysis patients during a five-Year observation period.', *Nutrients*, 15(5). Available at: <https://doi.org/10.3390/nu15051237>.
- Kalantar-Zadeh, K., Ficociello, L.H., Bazzanella, J., Mullon, C., et al. (2021) 'Slipping through the pores: hypoalbuminemia and albumin loss during hemodialysis.', *International journal of nephrology and renovascular disease*, 14, pp. 11–21.
- Kang, S.H., Kim, A.Y., Kim, J.C. and Do, J.Y. (2021) 'Comparison of body composition, strength, and physical performance measurements between healthy participants and hemodialysis patients.', *International journal of general medicine*, 14, pp. 7173–7179.
- KDOQI (2020) *KDOQI Clinical Practice Guideline for Nutrition in Chronic Kidney Disease: 2020 Update*.
- Keane, D., Gardiner, C., Lindley, E., Lines, S., et al. (2016) 'Changes in body composition in the two years after initiation of haemodialysis: a retrospective cohort study', *Nutrients*, 8(11), p. 702.
- Kim, J.-K., Choi, S.R., Choi, M.J., Kim, S.G., et al. (2014) 'Prevalence of and factors associated with sarcopenia in elderly patients with end-stage renal disease.', *Clinical nutrition (Edinburgh, Scotland)*, 33(1), pp. 64–68.
- Knuth, C.M., Auger, C. and Jeschke, M.G. (2021) 'Burn-induced hypermetabolism and skeletal muscle dysfunction.', *American journal of physiology. Cell physiology*, 321(1), pp. C58–C71.
- Kovesdy, C.P., Matsushita, K., Sang, Y., Brunskill, N.J., et al. (2018) 'Serum potassium and adverse outcomes across the range of kidney function: a CKD Prognosis Consortium meta-analysis.', *European heart journal*, 39(17), pp. 1535–1542. Available at: <https://doi.org/10.1093/eurheartj/ehy100>.
- Kraut, J.A. and Madias, N.E. (2017) 'Adverse effects of the metabolic acidosis of chronic kidney disease.', *Advances in chronic kidney disease*, 24(5), pp. 289–

297.

- Kuma, A. and Kato, A. (2022) 'Lifestyle-related risk factors for the incidence and progression of chronic kidney disease in the healthy young and middle-aged population', *Nutrients*, 14(18), p. 3787.
- Kurajoh, M., Mori, K., Miyabe, M., Matsufuji, S., et al. (2022) 'Nutritional status association with sarcopenia in patients undergoing maintenance hemodialysis assessed by nutritional risk index.', *Frontiers in nutrition*, 9, p. 896427.
- Larsson, L., Degens, H., Li, M., Salviati, L., et al. (2019) 'Sarcopenia: aging-related loss of muscle mass and function.', *Physiological reviews*, 99(1), pp. 427–511. Available at: <https://doi.org/10.1152/physrev.00061.2017>.
- Lee, M.-K., Jeong, H.H., Kim, M.-J., Ryu, H., et al. (2022) 'Nutrients against Glucocorticoid-Induced Muscle Atrophy.', *Foods (Basel, Switzerland)*, 11(5). Available at: <https://doi.org/10.3390/foods11050687>.
- Lemeshow, S., Jr, D.W.H., Klar, J. and Lwanga, S.K. (1997) *Besar Sampel Dalam Penelitian Kesehatan*. Yogyakarta: Gadjah Mada University Press.
- Lim, H.-S., Kim, H.-S., Kim, J.K., Park, M., et al. (2019) 'Nutritional status and dietary management according to hemodialysis duration', *Clinical nutrition research*. Department of Food Sciences & Nutrition, Yeonsung University, Anyang 14011, Korea., pp. 28–35.
- Lin, T.-Y., Peng, C.-H., Hung, S.-C. and Tarng, D.-C. (2018) 'Body composition is associated with clinical outcomes in patients with non-dialysis-dependent chronic kidney disease', *Kidney International*, 93(3), pp. 733–740.
- Lv, X., Zhou, C., Yan, Q., Tan, Z., et al. (2022) 'Elucidating the underlying mechanism of amino acids to regulate muscle protein synthesis: Effect on human health', *Nutrition*, 103–104, p. 111797.
- Mahan, L.K. and Raymond, J. (2016) *Krause's food & the nutrition care process*. 14th edn. St. Louis.
- Marcelli, D., Brand, K., Ponce, P., Milkowski, A., et al. (2016) 'Longitudinal changes in body composition in patients after initiation of hemodialysis therapy: results from an international cohort', *Journal of Renal Nutrition*, 26(2), pp. 72–80.
- Martinson, M., Ikizler, T.A., Morrell, G., Wei, G., et al. (2014) 'Associations of

- body size and body composition with functional ability and quality of life in hemodialysis patients.', *Clinical journal of the American Society of Nephrology: CJASN*, 9(6), pp. 1082–1090.
- Mathilakath, N.C., Selvaraj, J., Parameswaran, S., Viswanathan, S., et al. (2022) 'Prevalence of overhydration in patients on maintenance haemodialysis as determined by Body Composition Monitor and effects of attaining target dry weight.', *Cureus*, 14(9), p. e29509.
- McCance, K. and Huether, S. (2019) *Pathophysiology: The Biology Basis for Disease in Adults and Children*. 8th edn. Edited by Elsevier. Missouri.
- Di Micco, R., Krizhanovsky, V., Baker, D. and d'Adda di Fagagna, F. (2021) 'Cellular senescence in ageing: from mechanisms to therapeutic opportunities', *Nature Reviews Molecular Cell Biology*, 22(2), pp. 75–95.
- Moberg, M., Apró, W., Ekbom, B., Van Hall, G., et al. (2016) 'Activation of mTORC1 by leucine is potentiated by branched-chain amino acids and even more so by essential amino acids following resistance exercise', *American Journal of Physiology-Cell Physiology*, 310(11), pp. C874–C884.
- Molina, P., Vizcaíno, B., Molina, M.D., Beltrán, S., et al. (2018) 'The effect of high-volume online haemodiafiltration on nutritional status and body composition: the ProtEin Stores prEservaTion (PESET) study.', *Nephrology, dialysis, transplantation: official publication of the European Dialysis and Transplant Association - European Renal Association*, 33(7), pp. 1223–1235.
- Montaseri, M., Dindarloo, M.M., Espahbodi, F. and Yazdani-Charati, J. (2022) 'Factors Influencing Hemodialysis Patients Survival Using Frailty Models'.
- Moore, D.R., Churchward-Venne, T.A., Witard, O., Breen, L., et al. (2015) 'Protein ingestion to stimulate myofibrillar protein synthesis requires greater relative protein intakes in healthy older versus younger men.', *The journals of gerontology. Series A, Biological sciences and medical sciences*, 70(1), pp. 57–62. Available at: <https://doi.org/10.1093/gerona/glu103>.
- Morton, R.W., Taylor, D.A., Weijs, P.J.M. and Phillips, S.M. (2018) 'Defining anabolic resistance: implications for delivery of clinical care nutrition.', *Current opinion in critical care*, 24(2), pp. 124–130.
- Moser, S.E., Brown, A.M., Clark, B.C., Arnold, W.D., et al. (2022) 'Neuromuscular

- mechanisms of weakness in a mouse model of chronic alcoholic myopathy', *Alcoholism: Clinical and Experimental Research*, 46(9), pp. 1636–1647.
- Moustakim, R., Mziwira, M., El Ayachi, M. and Belahsen, R. (2020) 'Assessment of nutritional status, dietary intake and adherence to dietary recommendations in hemodialysis patients', *GSC Advanced Research and Reviews*, 3(2 SE-Original Article), pp. 9–19.
- Neugarten, J. and Golestaneh, L. (2019) 'Influence of sex on the progression of chronic kidney disease', in *Mayo Clinic Proceedings*. Elsevier, pp. 1339–1356.
- Nieves-Anaya, I., Vargas, M.B., Mayorga, H., García, O.P., et al. (2021) 'Comparison of nutritional and hydration status in patients undergoing twice and thrice-weekly hemodialysis: a silent drama in developing countries.', *International urology and nephrology*, 53(3), pp. 571–581.
- Nogami, E., Miyai, N., Zhang, Y., Sakaguchi, M., et al. (2021) 'Association of cigarette smoking with muscle mass reduction and Low muscle strength in community-dwelling elderly men', *Nihon Eiseigaku zasshi. Japanese Journal of Hygiene*, 76.
- Peng, Y., Fang, Z., Liu, M., Wang, Z., et al. (2019) 'Testosterone induces renal tubular epithelial cell death through the HIF-1 $\alpha$ /BNIP3 pathway', *Journal of translational medicine*, 17, pp. 1–13.
- PERNEFRI (2016) *9th Report of Indonesian Renal Registry*.
- PERNEFRI (2011) *Konsensus Nutrisi pada Penyakit Ginjal Kronis*.
- PERNEFRI (2020) *13th Annual Report of Indonesian Renal Registry*. Bandung.
- Post, A., Schutten, J.C., Kremer, D., van der Veen, Y., et al. (2021) 'Creatine homeostasis and protein energy wasting in hemodialysis patients', *Journal of Translational Medicine*, 19(1), p. 115.
- Rahayu, F., Fernandoz, T. and Ramlis, R. (2018) 'Hubungan frekuensi hemodialisis dengan tingkat stres pada pasien gagal ginjal kronik yang menjalani hemodialisis', *Jurnal keperawatan silampari*, 1(2), pp. 139–153.
- Raphael, K.L. (2019) 'Metabolic acidosis in CKD: core curriculum 2019.', *American journal of kidney diseases: the official journal of the National Kidney Foundation*, 74(2), pp. 263–275.

- Reid-McCann, R.J., Brennan, S.F., McKinley, M.C. and McEvoy, C.T. (2022) ‘The effect of animal versus plant protein on muscle mass, muscle strength, physical performance and sarcopenia in adults: protocol for a systematic review.’, *Systematic reviews*, 11(1), p. 64.
- Riani, A.P. (2019) ‘Hubungan asupan enegi dan protein dengan status gizi berdasarkan %LILA menurut umur pada pasien Chronic Kidney Disease on Hemodialysis di RSUD Dr. Saiful Anwar Malang’, *Labora Medika*, 3(1).
- Rokhmah, U.F., Purnamasari, D.U. and Saryono (2017) ‘Faktor-faktor yang berhubungan dengan penurunan nafsu makan pada pasien gagal ginjal kronik yang menjalani terapi hemodialisis (Studi kasus di RSUD Prof. Dr. Margono Soekarjo)’, *Jurnal Gizi Dan Pangan Soedirman*, 1(01), pp. 23–35.
- Rondanelli, M., Nichetti, M., Peroni, G., Faliva, M.A., et al. (2020) ‘Where to Find Leucine in Food and How to Feed Elderly With Sarcopenia in Order to Counteract Loss of Muscle Mass: Practical Advice.’, *Frontiers in nutrition*, 7, p. 622391. Available at: <https://doi.org/10.3389/fnut.2020.622391>.
- Rosa-Caldwell, M.E. and Greene, N.P. (2019) ‘Muscle metabolism and atrophy: let’s talk about sex.’, *Biology of sex differences*, 10(1), p. 43.
- Sabatino, A., Cuppari, L., Stenvinkel, P., Lindholm, B., et al. (2021) ‘Sarcopenia in chronic kidney disease: what have we learned so far?’, *Journal of nephrology*, 34(4), pp. 1347–1372.
- Sahathevan, S., Se, C.H., Ng, S.H., Chinna, K., et al. (2015) ‘Assessing protein energy wasting in a Malaysian haemodialysis population using self-reported appetite rating: a cross-sectional study.’, *BMC nephrology*, 16, p. 99.
- Sahathevan, S., Khor, B.-H., Ng, H.-M., Gafor, A.H.A., et al. (2020) ‘Understanding development of malnutrition in hemodialysis patients: A narrative review.’, *Nutrients*, 12(10).
- Salamah, S., Post, A., Alkaff, F.F., van Vliet, I.M.Y., et al. (2024) ‘Association between objectively measured protein intake and muscle status, health-related quality of life, and mortality in hemodialysis patients’, *Clinical Nutrition ESPEN* [Preprint].
- Seto, Y., Kimura, M., Matsunaga, T., Miyasita, E., et al. (2022) ‘Long-term body composition changes in patients undergoing hemodialysis: a single-center

- retrospective study', *Renal Replacement Therapy*, 8(1), p. 57.
- Shaaker, H. and Davenport, A. (2024) 'Does Dietary Intake Differ in Kidney Failure Patients With Sarcopenia and Frailty Treated by Hemodialysis', *Gerontology and Geriatric Medicine*, 10, p. 23337214241253430.
- Sinusi, R. and Hargono, A. (2021) 'Diabetes, hypertension, obesity, and smoking as risk factors for chronic kidney disease in productive age', *Jurnal Berkala Epidemiologi (JBE)*, 9(1), pp. 88–95.
- Siregar, C. (2020) *Buku Ajar Manajemen Komplikasi Pasien Hemodialisa*. Sleman: Deepublish.
- Song, Y., March, D.S., Biruete, A., Kistler, B.M., et al. (2022) 'A comparison of dietary intake between individuals undergoing maintenance hemodialysis in the United Kingdom and China.', *Journal of renal nutrition: the official journal of the Council on Renal Nutrition of the National Kidney Foundation*, 32(2), pp. 224–233. Available at: <https://doi.org/10.1053/j.jrn.2021.03.003>.
- Stack, A.G., Yermak, D., Roche, D.G., Ferguson, J.P., et al. (2016) 'Differential impact of smoking on mortality and kidney transplantation among adult Men and Women undergoing dialysis', *BMC nephrology*, 17, pp. 1–12.
- Stokes, T., Hector, A.J., Morton, R.W., McGlory, C., et al. (2018) 'Recent Perspectives Regarding the Role of Dietary Protein for the Promotion of Muscle Hypertrophy with Resistance Exercise Training.', *Nutrients*, 10(2). Available at: <https://doi.org/10.3390/nu10020180>.
- Sugiyono (2017) *Statistika untuk Penelitian*. Bandung: Penerbit Alfabeta.
- Sultan, S., Nasir, K., Qureshi, R., Dhrolia, M., et al. (2021) 'Assessment of the nutritional status of the hemodialysis patients by anthropometric measurements.', *Cureus*, 13(10), p. e18605.
- Sumetri (2019) 'Hubungan lama menjalani hemodialisis dengan asupan zat besi, protein dan kadar hemoglobin pada pasien hemodialisis di RSUD dr. M. Yunus Bengkulu'. Universitas Jenderal Soedirman.
- Surialaga, S. and IbnuSantosa, R.G. (2018) 'Gambaran persentase total massa otot dan total massa lemak tubuh pada golongan dewasa muda', *Prosiding Pendidikan Dokter*, pp. 613–619.
- Suryani, I., Isdiany, N. and Kusumayanti, G. (2018) *Dietetik Penyakit Tidak*

- Menular*. Jakarta: Kementerian Kesehatan RI.
- Tamam, B., Syah, D., Suhartono, M.T., Kusuma, W.A., et al. (2019) ‘Proteomic study of bioactive peptides from tempe’, *Journal of Bioscience and Bioengineering*, 128(2), pp. 241–248.
- Thompson, A.E., Anisimowicz, Y., Miedema, B., Hogg, W., et al. (2016) ‘The influence of gender and other patient characteristics on health care-seeking behaviour: a QUALICOPC study’, *BMC family practice*, 17, pp. 1–7.
- Vliet, S., Skinner, S.K., Beals, J.W., Pagni, B.A., et al. (2018) ‘Dysregulated Handling of Dietary Protein and Muscle Protein Synthesis After Mixed-Meal Ingestion in Maintenance Hemodialysis Patients’, *Kidney International Reports*, 3(6), pp. 1403–1415.
- Wang, J., Xu, M.-C., Huang, L.-J., Li, B., et al. (2023) ‘Value of neutrophil-to-lymphocyte ratio for diagnosing sarcopenia in patients undergoing maintenance hemodialysis and efficacy of Baduanjin exercise combined with nutritional support.’, *Frontiers in neurology*, 14, p. 1072986.
- Wijayanti, T.S. (2015) *Hubungan Asupan Energi dan Protein dengan Status Gizi Pasien Gagal Ginjal Kronik yang Menjalani Haemodialisis*. Universitas Jenderal Soedirman.
- Wisdom, K.M., Delp, S.L. and Kuhl, E. (2015) ‘Use it or lose it: multiscale skeletal muscle adaptation to mechanical stimuli’, *Biomechanics and Modeling in Mechanobiology*, 14(2), pp. 195–215.
- Wu, G. (2016) ‘Dietary protein intake and human health’, *Food Funct.*, 7(3), pp. 1251–1265. Available at: <https://doi.org/10.1039/C5FO01530H>.
- Yamamoto, S., Bieber, B.A., Komaba, H., Hanafusa, N., et al. (2022) ‘Medical director practice of advising increased dietary protein intake in hemodialysis patients with hyperphosphatemia: associations with mortality in the dialysis outcomes and practice patterns study’, *Journal of Renal Nutrition*, 32(2), pp. 243–250.
- Zha, Y. and Qian, Q. (2017) ‘Protein nutrition and malnutrition in CKD and ESRD.’, *Nutrients*, 9(3). Available at: <https://doi.org/10.3390/nu9030208>.