

DAFTAR PUSTAKA

- Amiryousefi, A., Hyvönen, J., & Poczai, P. (2018). iMEC: Online Marker Efficiency Calculator. *Applications in Plant Sciences*, 6(6), 4–7. <https://doi.org/10.1002/aps3.1159>
- Balada, C., Castro, M., Fassio, C., Zamora, A., Marchant, M. J., Acevedo, W., & Guzmán, L. (2021). Genetic diversity and biological activity of *Curcuma longa* ecotypes from Rapa Nui using molecular markers. *Saudi Journal of Biological Sciences*, 28(1), 707–716. <https://doi.org/10.1016/j.sjbs.2020.10.062>
- Baskaran, V., Jayanthi, R., Janakiram, T., & Abirami, K. (2009). Studies on Genetic Variability, Heritability and Genetic Advance in *Chrysanthemum*. *Journal of Horticultural Sciences*, 4(2), 174–176. <https://doi.org/10.24154/jhs.v4i2.540>
- Botstein, D., White, R. L., Skolnick, M., & Davis, R. W. (1980). Botstein. *Am J Hum Gen*, 32, 314–331. [papers2://publication/uuid/0B80518E-A22B-41F3-BE43-171F51007E42](https://pubmed.ncbi.nlm.nih.gov/601312/)
- BPS. (2022). *Produksi Tanaman Biofarmaka (Obat)*. <https://www.bps.go.id/id/statistics-table/2/NjMjMg==/produksi-tanaman-biofarmaka--obat-.html>
- Chalbi, A., Chikh-Rouhou, H., Mezghani, N., Slim, A., Fayos, O., Bel-Kadhi, M. S., & Garcés-Claver, A. (2023). Genetic Diversity Analysis of Onion (*Allium cepa* L.) from the Arid Region of Tunisia Using Phenotypic Traits and SSR Markers. *Horticulturae*, 9(10), 1–16. <https://doi.org/10.3390/horticulturae9101098>
- Choudhary, A. K., & Rahi, S. (2018). Organic cultivation of high yielding turmeric (*Curcuma longa* L.) cultivars: a viable alternative to enhance rhizome productivity, profitability, quality and resource-use efficiency in monkey–menace areas of north-western Himalayas. *Industrial Crops and Products*, 124(June), 495–504. <https://doi.org/10.1016/j.indcrop.2018.07.069>
- Doyle, JJ; Doyle, J. (1990). Isolation of Plant DNA From Fresh Tissue Plant. In *Focus* (Vol. 12, Issue 1, pp. 13–15).
- Evanno, G., Regnaut, S., & Goudet, J. (2005). Detecting the number of clusters of individuals using the software STRUCTURE: A simulation study. *Molecular Ecology*, 14(8), 2611–2620. <https://doi.org/10.1111/j.1365-294X.2005.02553.x>
- Ghelichpour, M., Shabani, A., & Shabanpour, B. (2013). Microsatellite variations and genetic structure of common carp (*Cyprinus carpio*) populations in Gomishan bay and Gorganroud River (Southeast of the Caspian Sea). *International Journal of Aquatic Biology*, 2013(1), 22–27.
- Govindaraj, M., Vetriventhan, M., & Srinivasan, M. (2015). Importance of genetic diversity assessment in crop plants and its recent advances: An overview of its analytical perspectives. *Genetics Research International*, 2015(Figure 1). <https://doi.org/10.1155/2015/431487>

- Jacqueline Batley. (2015). Plant Genotyping: Methods and Protocols. In *Plant Genotyping: Methods and Protocols* (Vol. 1245, pp. 77–89). Springer Science+Business Media New York. <https://doi.org/10.1007/978-1-4939-1966-6>
- Jan, H. U., Rabbani, M. A., & Shinwari, Z. K. (2011). Assessment of genetic diversity of indigenous turmeric (*Curcuma longa* L.) germplasm from Pakistan using RAPD markers. *Journal of Medicinal Plants Research*, 5(5), 823–830.
- Kaewkroek, K., Wattanapiromsakul, C., & Tewtrakul, S. (2009). Nitric oxide inhibitory substances from *Curcuma mangga* rhizomes. *Songklanakarin Journal of Science and Technology*, 31(3), 293–297.
- Kanaka, K. K., Sukhija, N., Goli, R. C., Singh, S., Ganguly, I., Dixit, S. P., Dash, A., & Malik, A. A. (2023). On the concepts and measures of diversity in the genomics era. *Current Plant Biology*, 33(March), 100278. <https://doi.org/10.1016/j.cpb.2023.100278>
- Karthikeyan, A., Young, K. N., Moniruzzaman, M., Beyene, A. M., Do, K., Kalaiselvi, S., & Min, T. (2021). Curcumin and its modified formulations on inflammatory bowel disease (Ibd): The story so far and future outlook. *Pharmaceutics*, 13(4), 1–38. <https://doi.org/10.3390/pharmaceutics13040484>
- Kumar, A., & Kaur, V. (2010). Characterisation and Evaluation of PGR: Principles and Techniques. *Division of Germplasm Evaluation, ICAR-NBPGR, New Delhi, August*, 139–145.
- Kumar, P., Dolkar, R., Manjunatha, G., & Pallavi, H. M. (2017). Molecular fingerprinting and assessment of genetic variations among advanced breeding lines of *Moringa oleifera* L. by using seed protein, RAPD and Cytochrome P 450 based markers. *South African Journal of Botany*, 111(May), 60–67. <https://doi.org/10.1016/j.sajb.2017.03.024>
- Li, Y. L., & Liu, J. X. (2018). StructureSelector: A web-based software to select and visualize the optimal number of clusters using multiple methods. *Molecular Ecology Resources*, 18(1), 176–177. <https://doi.org/10.1111/1755-0998.12719>
- Liu, B. H. (1998). *Statistical genomics: Linkage, mapping and QTL analysis*. CRC Press, Boca Raton, Florida, USA.
- Mallikarjuna, K. N., Tomar, B. S., Mangal, M., Singh, N., Singh, D., Kumar, S., Tomer, A., Singh, B., & Jat, G. S. (2023). Genetic Diversity and Population Structure Analyses in Bitter Gourd (*Momordica charantia* L.) Based on Agro-Morphological and Microsatellite Markers. *Plants*, 12(19). <https://doi.org/10.3390/plants12193512>
- Mbhele, Z., Zharare, G. E., Zimudzi, C., & Ntuli, N. R. (2023). Assessing Genetic Variation among *Strychnos spinosa* Lam. Morphotypes Using Simple Sequence Repeat Markers. *Plants*, 12(15), 1–13. <https://doi.org/10.3390/plants12152810>
- Nandakumar, K., Fakrudin, B., V, M. K. N., Brijesh, A., & Venkatesha, J. (2023). Genetic Diversity in Turmeric (*Curcuma longa* L.) Genotypes using Molecular Markers. 15(2), 1229–1236.

- Panwar, P., & Netrapal, R. K. S. (2010). *Efficiency of RAPD , SSR and Cytochrome P 450 gene based markers in accessing genetic variability amongst finger millet (Eleusine coracana) accessions*. 4075–4082. <https://doi.org/10.1007/s11033-010-0067-5>
- Peakall, R., & Smouse, P. E. (2012). GenALEx 6.5: Genetic analysis in Excel. Population genetic software for teaching and research-an update. *Bioinformatics*, 28(19), 2537–2539. <https://doi.org/10.1093/bioinformatics/bts460>
- Powell, W., Machray, G. C., & Proven, J. (1996). Polymorphism revealed by simple sequence repeats. *Trends in Plant Science*, 1(7), 215–222. [https://doi.org/10.1016/1360-1385\(96\)86898-1](https://doi.org/10.1016/1360-1385(96)86898-1)
- Prasad S & Aggarwal BB. (2011). *Turmeric, the golden spice, Herbal Medicine: Biomolecular and Clinical Aspects* (2nd editio). CRC Press/Taylor & Francis, Boca Raton (FL).
- Prasetia, R. G. N., Suprayogi, Asnani, A., Oktaviani, E., & Nuryana, I. (2022). Microsatellite Markers and Metabolite Profiles of Salt-Tolerant Rice: Inpari Unsoed 79 Agritan. *Caraka Tani: Journal of Sustainable Agriculture*, 37(2), 223–232. <https://doi.org/10.20961/carakatani.v37i2.57728>
- Prevost, A., & Wilkinson, M. J. (1999). A new system of comparing PCR primers applied to ISSR fingerprinting of potato cultivars. *Theoretical and Applied Genetics*, 98(1), 107–112. <https://doi.org/10.1007/s001220051046>
- Pritchard JK, Stephens M, D. P. (2008). Inference of Population Structure Using Multilocus Genotype Data. *Genetics* =, 164(2), 593–601.
- Pritchard JK, W. (2003). *Documentation for the structure software, version 2*. Department of Human Genetics, University of Chicago.
- Putri, T. K., Anindita, P. A., Wicaksana, N., Suganda, T., Concibido, V., Karuniawan, A., Tenggara, A., & Aggarwal, P. (2020). Genetic Diversity of 64 Turmeric Accessions from Indonesia Based on P450- Based Analogue (PBA). *Buletin Penelitian Tanaman Rempah Dan Obat*, 31(2), 123–134.
- Rajkumari, S., & Sanatombi, K. (2018). Nutritional value, phytochemical composition, and biological activities of edible Curcuma species: A review. *International Journal of Food Properties*, 20(3), S2668–S2687. <https://doi.org/10.1080/10942912.2017.1387556>
- Ravi, R. S. D., & Bindu, E. A. S. (2020). The efficiency of Cytochrome P450 gene - based markers in accessing genetic variability of drumstick (Moringa oleifera Lam .) accessions. *Molecular Biology Reports*, 0123456789. <https://doi.org/10.1007/s11033-020-05391-w>
- Rocha, S. S., Londe, L. C. N., Pimenta, S., Cardoso, M. M., Gonçalves, N. P., Gomes, W. S., & Calaes, J. G. (2020). Congruence between morphological and molecular markers for genetic diversity analysis applied to forage palm genotypes propagated via bioreactors. *Industrial Crops and Products*, 147(February), 112230. <https://doi.org/10.1016/j.indcrop.2020.112230>
- Royal Botanic Gardens Kew. (2021). *Curcuma Longa* L. available:

- Salgotra, R. K. . C., & B.S. (2023). Genetic Diversity, Conservation, and Utilization of Plant Genetic Resources. *Gene*, 14, 174. <https://doi.org/10.3390/genes14010174>
- Salgotra, R. K., Gupta, B. B., Bhat, J. A., & Sharma, S. (2015). Genetic diversity and population structure of basmati rice (*Oryza sativa* L.) Germplasm collected from North Western Himalayas using trait linked SSR markers. *PLoS ONE*, 10(7), 1–19. <https://doi.org/10.1371/journal.pone.0131858>
- Sasikumar, B. (2005). Genetic resources of curcuma diversity, characterization and utilization. *Plant Gen, Res.*, 3(2), 230–251.
- Schalk, M., Nedelkina, S., Schoch, G., Batard, Y., & Werck-Reichhart, D. (1999). Role of unusual amino acid residues in the proximal and distal heme regions of a plant P450, CYP73A1. *Biochemistry*, 38(19), 6093–6103. <https://doi.org/10.1021/bi982989w>
- Shen, J. ling, Jia, X. nan, Ni, H. qun, Sun, P. guang, Niu, S. hui, & Chen, X. yang. (2010). AFLP analysis of genetic diversity of jatropha curcas grown in Hainan, China. *Trees - Structure and Function*, 24(3), 455–462. <https://doi.org/10.1007/s00468-010-0413-1>
- Singh, T. J., Patel, R. K., kumar, S., Patel, N., & Patel, P. A. (2018). Molecular Diversity Analysis in Turmeric (*Curcuma longa* L.) Using SSR Markers. *International Journal of Current Microbiology and Applied Sciences*, 7(11), 552–560. <https://doi.org/10.20546/ijcmas.2018.711.066>
- Soudamini, K. K., and Kuttan, R. (1989). Drugs and chemicals Dried turmeric tubers were purchased from a local market (authenti- cated by CD. Varghese, Botanist, St. Thomas College, Trichurl and. *J. Ethnopharmacol* 27, 27, 227–233.
- Sunaryo, W., Wahida, Idris, S. D., Pratama, A. N., Ratanasut, K., & Nurhasanah. (2020). Genetic relationships among cultivated and wild bananas from east Kalimantan, Indonesia based on ISSR markers. *Biodiversitas*, 21(2), 824–832. <https://doi.org/10.13057/biodiv/d210250>
- Tanksley, S. D., & McCouch, S. R. (1997). Seed banks and molecular maps: Unlocking genetic potential from the wild. *Science*, 277(5329), 1063–1066. <https://doi.org/10.1126/science.277.5329.1063>
- Tessier, C., David, J., This, P., Boursiquot, J. M., & Charrier, A. (1999). Optimization of the choice of molecular markers for varietal identification in *Vitis vinifera* L. *Theoretical and Applied Genetics*, 98(1), 171–177. <https://doi.org/10.1007/s001220051054>
- Varshney, R. K., Graner, A., & Sorrells, M. E. (2005). Genic microsatellite markers in plants: Features and applications. *Trends in Biotechnology*, 23(1), 48–55. <https://doi.org/10.1016/j.tibtech.2004.11.005>
- Verma, S., Singh, S., Sharma, S., Tewari, S. K., Roy, R. K., Goel, A. K., & Rana, T. S. (2015). Assessment of genetic diversity in indigenous turmeric (*Curcuma longa*) germplasm from India using molecular markers. *Physiology and Molecular Biology of Plants*, 21(2), 233–

242. <https://doi.org/10.1007/s12298-015-0286-2>

- Wicaksana, N., Gilani, S. A., Ahmad, D., Kikuchi, A., & Watanabe, K. N. (2011). Morphological and molecular characterization of underutilized medicinal wild ginger (*Zingiber barbatum* Wall.) from Myanmar. *Plant Genetic Resources: Characterisation and Utilisation*, 9(4), 531–542. <https://doi.org/10.1017/S1479262111000840>
- Winter, P., & Kahl, G. (1995). Molecular marker technologies for plant improvement. *World Journal of Microbiology & Biotechnology*, 11(4), 438–448. <https://doi.org/10.1007/BF00364619>
- Wright, S. (1978). *Evolution and the Genetics of Populations : Variability Within and Among Natural Populations*. University of Chicago Press.
- Yamanaka, S., Suzuki, E., Tanaka, M., Takeda, Y., Watanabe, J. A., & Watanabe, K. N. (2003). Assessment of cytochrome P450 sequences offers a useful tool for determining genetic diversity in higher plant species. *Theoretical and Applied Genetics*, 108(1), 1–9. <https://doi.org/10.1007/s00122-003-1403-0>
- Zatybekov, A., Yermagambetova, M., Genievskaya, Y., Didorenko, S., & Abugalieva, S. (2023). Genetic Diversity Analysis of Soybean Collection Using Simple Sequence Repeat Markers. *Plants*, 12(19). <https://doi.org/10.3390/plants12193445>
- Záveská, E., Fér, T., Šída, O., Krak, K., Marhold, K., & Leong-Skornickova, J. (2012). Phylogeny of *Curcuma* (Zingiberaceae) based on plastid and nuclear sequences: Proposal of the new subgenus *Ecomata*. *Taxon*, 61(4), 747–763. <https://doi.org/10.1002/tax.614004>