

DAFTAR PUSTAKA

- Abebe, E. & Coomans, A. 1996. Aquatic nematodes from Ethiopia VI. *Hydrobiologia*, 332: 41–6. <https://doi.org/10.1007/BF00020779>
- Abo-Korah, M. S., M. E. Sweelam, & A. M. Yassin. 2022. Effect of different bioagents on the population density of meloidogyne incognita infected tomato plants. *Egyptian Journal of Agronematology*, 21(1):14–22
- Altash, S., Kostadinova, A., & Peneva, V. 2024. Integrative taxonomic study of mononchid nematodes from riparian habitats in Bulgaria. I. Genera *Mononchus* Bastian, 1865 and *Coomansus* Jairajpuri & Khan, 1977 with the description of *Mononchus pseudoaquaticus* sp. nov. and a key to the species of *Mononchus*. *ZooKeys*, 1206: 137.
- Amir, F., Widajati, W., Rahmadhini, N., & Imanadi, L. 2025. Nematoda yang berasosiasi dengan tanaman padi (*Oryza Sativa* L. di Desa Sumberngepoh, Lawang, Kabupaten Malang. *Jurnal Agrotek Tropika*, 12(4): 757-768.
- Amulu, L. U., Oyedele, D. J., & Adekunle, O. K. 2021. Distribution of plant-parasitic nematodes and comparative effects of fadama (hydromorphic) fields and upland fields on plant-parasitic nematodes and free-living nematodes on three indigenous vegetables commonly grown in Osun State Nigeria. *Archives of Phytopathology and Plant Protection*, 54 (15): 1131–1142. <https://doi.org/10.1080/03235408.2021.1890912>
- Andrássy, I. 2012. On the “Paractinolaimus” genus group (Nematoda: Actinolaimidae), with description of five new and two rare species of Egtitus Thorne, 1967. *Journal of Natural History*, 46(7): 453-494. doi:10.1080/00222933.2011.642415
- Asghari, R., Eskandari, A., Maafi, Z. T., Alvarez-Ortega, S., & Zhao, Z. Q. 2017. Morphological and molecular characterisation of new and known species of *Tripyla* Bastian, 1865 (Triplonchida: Tripylidae) from northern Iran, with phylogenetic relationships, compendium and identification key. *Nematology*, 19(1): 21-56.
- Asyiah, I. N., Imam, M., Ankardiansyah, Pandu., P., Aris, B., Dwi, N., & Laeli, N. 2024. Effect of liquid biofertilizer bre4 and organic fertilizer on soil nematode diversity in arabica coffee plantation: pengaruh pupuk hayati cair bre4 dan pupuk organik terhadap keragaman nematoda tanah pada lahan kopi arabika. *Jurnal Penelitian Pertanian Terapan*, 24(3): 321-328.
- Bajaj, H. K., & Kanwar, R. S. 2015. Biology and predatory attributes of a diplogasterid nematode, Khan, 2008. *Helminthologia*, 52(1): 50-57.
- Baweja, P., Kumar, S., & Kumar, G. 2020. Fertilizers and pesticides: Their impact

- on soil health and environment. *Soil health*, 1(1): 265-285.
- Beare, M.H., Reddy, M.V., Tian, G. & Srivastava, S.C. 1997. Agricultural intensification, soil biodiversity and agroecosystem function in the tropics: The role of decomposer biota. *Applied Soil Ecology*, 6(1): 87–108
- Biswal, D. 2022. Nematodes as ghosts of land use past: elucidating the roles of soil nematode community studies as indicators of soil health and land management practices. *Applied Biochemistry and Biotechnology*, 194(5): 2357-2417.
- Bokhorst, S., J. Asplund, P. Kardol, and D. A. Wardle 2015. Lichen physiological traits and growth forms affect communities of associated invertebrates. *Ecology*, 96(9): 2394-2407.
- Bristol, D., Hassan, K., Blankinship, J. C., & Nielsen, U. N. 2023. Responses of nematode abundances to increased and reduced rainfall under field conditions: A meta-analysis. *Ecosphere*, 14(1): 43-64.
- Cabos, R.Y.M., Wang, K.H., Sipes, B.S., Heller, W.P., Matsumoto, T.K., 2013. Detection of plant-parasitic nematode DNA in the gut of predatory and omnivorous nematodes. *Nematropica*, 43(1): 44–48.
- Cakmak, T., Kaydan, M., Gözel, Ç., Pena-Santiago, R., Abolafia, J., & Gözel, U. 2021. New records of free-living nematode species from Mount Ararat, Turkey, with SEM study of three remarkable species. *Biologia*, 76(1).
- Čerevková, A., Renčo, M., Miklisová, D., & Gömöryová, E. 2021. Soil nematode communities in managed and natural temperate forest. *Diversity*. 13(7):1-13
- Chen, X., Xue, W., Xue, J., Griffiths, B. S., & Liu, M. 2020. Contribution of bacterivorous nematodes to soil resistance and resilience under copper or heat stress. *Soil Ecology Letters*, 2 (1): 220-229.
- Coleman, DC and Wall, DH. 2015. Soil Fauna: Occurrence, Biodiversity, and Roles in Ecosystem Function. P:120-125. In: E.A Paul (Ed.). 2023, *Soil Fauna. Soil Microbiology, Ecology and Biochemistry (4th ed.)*. London, Academic Press. doi: 10.1016/b978-0-12-415955-6.00005-0.
- Coomans, A. V., & Raski, D. J. 1988. Two new species of *Prismatolaimus* de Man, 1880 (Nemata: Prismatolaimidae) in Southern Chile. *Journal of Nematology*, 20(2): 288.
- Cornejo-Condori, G., Lima-Medina, I., Bravo-Portocarrero, R., Barzola-Tito, K., & Casa-Coila, V. 2021. Nematodes associated with andean papaya (*Carica pubescens* L.) in Sandia, Puno, Peru. *Bioagro*, 33(3):191-202.
- Darmawan, A., Arminudin, A. T., & Rahmadani, E. 2024. Identifikasi nematoda di Perkebunan Kelapa Sawit PTPN V Sei Pagar. *Jurnal*

Agroteknologi, 15(1): 37-46.

- Devi, G. 2020. Nematodes as environmental indicator. *IntlJ Curr Res*, 12 (05): 11699-11705.
- Devi, G., & George, J. 2018. Predatory nematodes as bio-control agent against plant-parasitic nematode-A review. *Agricultural Reviews*, 39(1): 55-61.
- Dionfriski, A., Siregar, S. H., & Nurrachmi, I. 2021. Epipellic diatom community structure in the intertidal zone Mengkapan Waters, Sungai Apit District, Siak Regency. *Journal of Coastal and Ocean Sciences*, 2(3): 207-216.
- Di Montanara, A. C., Baldrighi, E., Franzo, A., Catani, L., Grassi, E., Sandulli, R., & Semprucci, F. 2022. Free-living nematodes research: State of the art, prospects, and future directions. A bibliometric analysis approach. *Ecological Informatics*, 72(1): 101891.
- Du Preez, G., Daneel, M., De Goede, R., Du Toit, M. J., Ferris, H., Fourie, H., ... & Schmidt, J. H. 2022. Nematode-based indices in soil ecology: Application, utility, and future directions. *Soil Biology and Biochemistry*, 169(1): 108640.
- Du Preez, G. C., Daneel, M. S., Wepener, V., & Fourie, H. 2018. Beneficial nematodes as bioindicators of ecosystem health in irrigated soils. *Applied Soil Ecology*, 132(1): 155-168.
- Ferris H, Bongers T, RGM de Goede. 2001. A framework for soil food web diagnostics: Extension of the nematode faunal analysis concept. *Appl Soil Ecol*, 18(1):13-29.
- Ferris, H., Cares, J. E., & Esteves, A. M. 2021. Effect of land use and seasonality on nematode faunal structure and ecosystem functions in the Caatinga dry forest. *European Journal of Soil Biology*, 103(1): 103296.
- Ferris, V. R., & Ferris, J. M. 1989. Why ecologists need systematics: importance of systematics to ecological research. *Journal of nematology*, 21(3): 308–314.
- Gade, R. B., & Hiware, C. J. 2017. Studies on the soil nematode diversity in relation to the soil parameters from sugarcane fields of Aurangabad District Maharashtra State, India. *International Journal for Research in Applied Science & Engineering Technology*, 5(1): 1632-1640.
- Gagarin, V. G. 2001. *Seinura informis* sp. n. from European Russia and a key to species of the genus *Seinura* Fuchs, 1931 (Nematoda: Aphelenchoididae). *Zoosystematica Rossica*, 9(1): 19-23.
- Garcia, N., Grenier, E., Buisson, A., & Folcher, L. 2022. Diversity of plant parasitic nematodes characterized from fields of the French national monitoring progme for the Columbia root-knot nematode. *Plos one*, 17(3): e0265070.

- Gattoni, K., Gendron, E. M. S., Borgmeier, A., McQueen, J. P., Mullin, P. G., Powers, K., ... & Porazinska, D. L. 2022. Context-dependent role of abiotic and biotic factors structuring nematode communities along two environmental gradients. *Molecular Ecology*, 31(14): 3903-3916.
- Grootaert, P. & Maertens, D., 1976. Cultivation and life cycle of *Mononchus aquaticus*. *Nematologica*. 22(2): 173–181.
- Gu, J., Maria, M., Liu L. & Pedram, M. 2020. Description of *Seinura italiensis* n. sp. (Tylenchomorpha: Aphelenchoididae) found in the medium soil imported from Italy. *J Nematol*, 52:1-11. doi: 10.21307/jofnem-2020-018.
- Guan, P., Zhang, X., Yu. J., Cheng, Y., Li, Q., Andriuzzi, W.S. & Liang W. 2018. Soil microbial food web channels associated with biological soil crusts in desertification restoration: The carbon flow from microbes to nematodes. *Soil Biol Biochem*, 116 (1): 82-90.
- Gusakov, V. A. & Gagarin, V. G. 2015. *Dorylaimus proximus* sp. n. (Nematoda, Dorylaimida) from freshwater bodies of Vietnam. *International Journal of Nematology*, 25(2): 117-121.
- Gusakov, V. A., Gagarin, V. G., Shkil, F. N., & Zlenko, D. V. 2022. *Eudorylaimus chukotkanus* sp. n. and *E. mylnikovi* sp. n. (Nematoda: Dorylaimida) from Lake El'gygytgyn (Chukotka, Russia). *Inland Water Biology*, 15(2): 97-106.
- Hadi, H., D. Chalil & Rahmanta. 2017. Pengaruh perubahan iklim terhadap produksi padi sawah di Provinsi Sumatera Utara. *Social Economic of Agriculture and Agribusiness*, 8(5): 1- 12
- Hanum, L., Windusari, Y., Setiawan, A., Hidayat, M. R., Adriansyah, F., Mubarak, A. A., & Pratama, R. 2018. *Morfologi dan Molekuler Padi Lokal Sumatera Selatan*. Palembang: NoerFikri.
- Harni, R. 2016. Prospek pengembangan bakteri endofit sebagai agens hayati pengendalian nematoda/the prospect of developing endophytic bacteria as nematodes biological control. *Perspektif: Review Penelitian Tanaman Industri*, 15(1): 31-49.
- Heidemann, K., Hennies, A., Schakowske, J., Blumenberg, L., Ruess, L., Scheu, S., & Maraun, M. 2014. Free-living nematodes as prey for higher trophic levels of forest soil food webs. *Oikos*, 123(10): 1199-1211.
- Henne, S., Sombke, A., & Schmidt-Rhaesa, A. 2017. Immunohistochemical analysis of the anterior nervous system of the free-living nematode *Plectus* spp. (Nematoda, Plectidae). *Zoomorphology*, 136(2): 175-190.
- Herawati, W.D. 2012. *Budidaya Padi*. Javalitera. Yogyakarta.
- Heyns, J. and Argo, A. D. 1969. Actinolaimoidea of South Africa (Nematoda:

Dorylaimidae. *Phytophylactica*, 1(34): 217-228.

- Hodda, M. 2022. Phylum Nematoda: Feeding habits for all valid genera using a new, universal scheme encompassing the entire phylum, with descriptions of morphological characteristics of the stoma, a key, and discussion of the evidence for trophic relationships. *Zootaxa*, 5114(1): 318-451.
- Holovachov, O. & Shoshin, A. 2014. 7.4 Order Triplonchida Cobb, 1919. P. 251-276. In A. Schmidt-Rhaesa, *Volume 2 Nematoda*. Dey Gruter, Berlin.
- Hoogen, Van Den., J., Geisen, S., Routh, D., Ferris, H., Traunspurger, W., Wardle, D. A., ... & Crowther, T. W. 2019. Soil nematode abundance and functional group composition at a global scale. *Nature*, 572(7768): 194-198.
- Hu, C., & Qi, Y. C. 2010. Abundance and diversity of soil nematodes as influenced by different types of organic manure. *Helminthologia*, 47 (1): 58-66.
- Iliev, I., & Ilieva, Z. 2016. Soil nematodes of family Plectidae (Nematoda, Plectida) in different plant communities in Strandzha Mountain (Bulgaria, Turkey). *Silva Balcanica*, 17(1): 71-108
- Iqbal S, Jones MGK. 2017. *Nematodes*. Elsevier, Amsterdam. doi: 10.1016/B978-0-12-948076.00061-7.
- Kanwar, R. S., Patil, J. A., & Yadav, S. 2021. Prospects of using predatory nematodes in biological control for plant parasitic nematodes-A review. *Biological Control*. 160(1): 104668. <https://doi.org/10.1016/j.biocontrol.2021.104668>
- Kanzaki, N., Ekino, T., & Masuya, H. 2019. *Seinura caverna* n. sp. (Tylenchomorpha: Aphelenchoididae), an androdioecious species isolated from bat guano in a calcareous cave. *Nematology*, 21(2): 207-225.
- Kanzaki, N., Ekino, T., Hamaguchi, K., & Takeuchi-Kaneko, Y. 2021. Three *Seinura* species from Japan with a description of *S. shigaensis* n. sp. (Tylenchomorpha: Aphelenchoididae). *Plos one*, 16(1): e0244653.
- Kazemi, E., Niknam, G., Jabbari, H., & Peña-Santiago, R. 2018. *Eudorylaimus kahaqensis* sp. n. (Nematoda: Dorylaimida: Qudsianematidae), an interesting new species from Iran. *Zootaxa*, 4369(1): 128.
- Kekelis P, Papatheodorou EM, Terpsidou E, Dimou M, Aschonitis V, Monokrousos N. 2022. The free-living nematodes as indicators of the soil quality in relation to the clay content, when coffee waste is applied. *Agronomy*, 12 (2702): 1-16.
- Kergunteuil, A., Campos-Herrera, R., Sánchez-Moreno, S., Vittoz, P., &

- Rasmann, S. 2016. The abundance, diversity, and metabolic footprint of soil nematodes is highest in high elevation alpine grasslands. *Frontiers in Ecology and Evolution*, 4(1): 84.
- Khusna, I. M., & Mariana, N. 2021. Sistem Pendukung Keputusan Pemilihan Bibit Padi Berkualitas Dengan Metode AHP Dan Topsis. *Jurnal Sisfokom (Sistem Informasi Dan Komputer)*, 10(2): 162-169.
- Kim, J., Yu, J. N., & Kim, T. 2020. First report of two plectus species (nematoda: Plectida) from Korea. *Animal Systematics, Evolution and Diversity*, 36(3): 251-257.
- Kitagami, Y., Obase, K., & Matsuda, Y. 2022. High-throughput sequencing and conventional morphotyping show different soil nematode assemblages but similar community responses to altitudinal gradients on Mt. Ibuki, Japan. *Pedobiologia*, 90 (1):150788.
- Korobushkin, D. I., Butenko, K. O., Gongalsky, K. B., Saifutdinov, R. A., & Zaitsev, A. S. 2019. Soil nematode communities in temperate rice-growing systems. *European Journal of Soil Biology*, 93(1): 103099.
- Kouser, Y., Shah, A. A., & Rasmann, S. 2021. The functional role and diversity of soil nematodes are stronger at high elevation in the lesser Himalayan Mountain ranges. *Ecology and Evolution*, 11(20): 13793-13804.
- Kudrin, A. A., Tsurikov, S. M., & Tiunov, A. V. 2015. Trophic position of microbivorous and predatory soil nematodes in a boreal forest as indicated by stable isotope analysis. *Soil Biology and Biochemistry*, 86(1): 193–200. doi:10.1016/j.soilbio.2015.03.017
- Kumar, N., Kumar, A., Jeena, N., Singh, R., & Singh, H. 2020. Factors influencing soil ecosystem and agricultural productivity at higher altitudes. *Microbiological Advancements for Higher Altitude Agro-Ecosystems & Sustainability*, 1(1): 55-70.
- Laasli, S. E., Lahlali, R., Dababat, A. A., Paulitz, T., Assouguem, A., Ullah, R., ... & Mokrini, F. 2024. Soil health in Moroccan olive agroecosystems: The bioindication role of terrestrial nematode dynamics. *Heliyon*, 10(23): 1-18
- Lazarova, S., Coyne, D., Rodriguez, M. G., Peteira, B., & Ciancio, A. 2021. Functional diversity of soil nematodes in relation to the impact of agriculture—a review. *Diversity*, 13(2): 64.
- Li Z, Chen X, Li J, Liao X, Li D, He X, Zhang W, Zhao J. 2022. Relationships between soil nematode communities and soil quality as affected by land-use type. *Forests*,13(10): 1658
- Li, Q., Liang, W., Zhang, X., & Mahamood, M. 2017. *Soil nematodes of grasslands in Northern China*. Academic Press.

- Lu, Q., Liu, T., Wang, N., Dou, Z., Wang, K., & Zuo, Y. 2020. A review of soil nematodes as biological indicators for the assessment of soil health. *Front. Agric. Sci. Eng.*, 7(1): 275-281.
- Magfiroh, N., Lapanjang, I. M., & Made, U. 2017. Pengaruh jarak tanam terhadap pertumbuhan dan hasil tanaman padi (*Oryza sativa* L.) pada pola jarak tanam yang berbeda dalam sistem tabela. *Agrotekbis: Jurnal Ilmu Pertanian (e-journal)*, 5(2): 212-221.
- Majdi, N., & Traunspurger, W. 2015. Free-living nematodes in the freshwater food web: a review. *Journal of nematology*, 47(1): 28–44.
- McSorley, R. 2012. Ecology of the dorylaimid omnivore genera *Aporcelaimellus*, *Eudorylaimus* and *Mesodorylaimus*. *Nematology*, 14(6): 645-663.
- Mulyadi. 2009. *Nematologi Pertanian*. Gajah Mada University Press, Yogyakarta.
- Musarrat, A. R., Shahina, F., Shah, A. A., Saba, R., & Feroza, K. 2016. Community analysis of plant parasitic and free living nematodes associated with rice and soybean plantation from Pakistan. *Applied Ecology and Environmental Research*, 14(5): 19-33.
- Mutala'liah, M. L., Manan, A., & Bayyinah, L. N. I. 2023. Abundance and diversity of terrestrial free-living nematodes in potato agroecosystem. *Nusantara Bioscience*, 15(1): 129-135.
- Mwamula, A. O., Kim, Y. S., & Lee, D. W. 2024. Morphological and molecular characterization of *Paractinolaimus uljinensis* n. sp. (Nematoda: Actinolaimidae) from Korea, with an updated compendium of the genus. *Journal of Nematology*, 56(1): 20240040-20240040.
- Neher, D. A. 2010. Ecology of plant and free-living nematodes in natural and agricultural soil. *Annual Review of Phytopathology*, 48(1): 371–394. doi:10.1146/annurev-phyto-073009-114439
- Ney, L., Franklin, D., Mahmud, K., Cabrera, M., Hancock, D., Habteselassie, M., and Newcomer, Q. 2018. Examining trophic-level nematode community structure and nitrogen mineralization to assess local effective microorganisms' role in nitrogen availability of swine effluent to forage crops. *Applied Soil Ecology*, 130: 209–218. doi: 10.1016/j.apsoil.2018.06.015.
- Nisa, R. U., Tantray, A. Y., Kouser, N., Allie, K. A., Wani, S. M., Alamri, S. A., ... & Shah, A. A. 2021. Influence of ecological and edaphic factors on biodiversity of soil nematodes. *Saudi Journal of Biological Sciences*, 28(5): 3049-3059.
- Nisa, R.U., Nisa, A.U., Hroobi, A.A., Shah, A., & Tantray, A.Y. 2022. Year-long assessment of soil nematode diversity and root inhibition-indicator nematode genera in rice fields. *Biology*, 11(1572):1-20

- Okada, H., & Harada, H. 2007. Effects of tillage and fertilizer on nematode communities in a Japanese soybean field. *Applied Soil Ecology*, 35(3): 582-598.
- Oktafiyanto, M. F., & Ibrahim, R. 2021. Keragaman dan kelimpahan nematoda secara horizontal dan vertikal pada beberapa tanaman sayur di Kabupaten Cianjur. *Jurnal Agro Wiralodra*, 4(1): 9–15.
- Olatunji, O., Gong, S., Tariq, A., Pan, K., Sun, X., Chen, W., Zhang, L., Dakhil, M., Huang, D., & Tan, X. 2018. The effect of phosphorus addition, soil moisture, and plant type on soil nematode abundance and community composition. *Journal of Soils and Sediments*, 19 (1): 1139-1150
- Öztürk, L. 2023. A field study on determination of nematode diversity in canola fields in Tekirdağ, Turkey. *Türk Doğa ve Fen Dergisi*. 12(1): 91-98.
- Panahi, H., Shokoohi, E., Vinciguera, M. T., Clausi, M., & Mashela, P. W. 2019. *Paractinolaimus persicus* sp. n.(Nematoda: Actinolaimidae) from Iran. *Zootaxa*, 4551(2): 244-250.
- Pedram, M., Niknam, G., Vinciguerra, M. T., Ye, W., & Robbins, R. T. 2010. Description of *Paractinolaimus decraemerae* sp. n. and redescription of *P. parietinus* Eroshenko, 1977 (Dorylaimida: Actinolaimidae) from northwest Iran. *International Journal of Nematology*, 20(2): 169-178.
- Pedram, M., Niknam, G., Vinciguerra, M. T., Ye, W., & Robbins, R. T. 2011. Morphological and molecular characterization of *Paractinolaimus sahandi* n. sp.(Nematoda: Actinolaimidae) from the Sahand Mountains in Iran. *Journal of Helminthology*, 85(3): 276-282.
- Porazinska, D. L., Bueno de Mesquita, C. P., Farrer, E. C., Spasojevic, M. J., Suding, K. N., & Schmidt, S. K. 2021. Nematode community diversity and function across an alpine landscape undergoing plant colonization of previously unvegetated soils. *Soil Biology and Biochemistry*, 161, 108380: 1-11
- Pothula, S.K., Grewal PS, Auge RM, Saxton AM, Bernard EC. 2019. Agricultural intensification and urbanization negatively impact soil nematode richness and abundance: A meta-analysis. *J Nematol*, 51 (1): 1-17. doi: 10.21307/jofnem-2019-011.
- Purwansyah, T. S., Rosanti, D., & Kartika, T. 2021. Morfometri beberapa varietas tanaman padi (*Oryza sativa* L.) di Kecamatan Pulau Rimau Banyuasin. *Indobiosains*, 28-38.
- Quénéhervé, P., & Chotte, J. L. 1996. Distribution of nematodes in vertisol aggregates under a permanent pasture in Martinique. *Applied Soil Ecology*, 4(3): 193-200
- Renčo, M., D’addabbo, T., Sasanelli, N., Papajova, I. 2007. The effect of five composts of different origin on the survival and reproduction of

Globodera rostochiensis. *Nematology*, 9(4): 537 – 543

- Renčo, M., Gömöryová, E., & Čerevková, A. 2020. The effect of soil type and ecosystems on the soil nematode and microbial communities. *Helminthologia*, 57(2): 129-144.
- Roidah, I. S. 2015. Analisis pendapatan usahatani padi musim hujan dan musim kemarau (studi kasus di Desa Sepatan Kecamatan Gondang Kabupaten Tulungagung). *Agribis*, 11(13): 45-55.
- Rosanti, D. 2016. Taksonomi gulma padi (*Oryza sativa*) di areal persawahan Jakabaring Palembang. *Sainmatika: Jurnal Ilmiah Matematika dan Ilmu Pengetahuan Alam*, 13(1): 46-51
- Rosanti, D. 2016. Taksonomi Gulma Padi (*Oryza sativa*) di Areal Persawahan Jakabaring Palembang. *Sainmatika: Jurnal Ilmiah Matematika dan Ilmu Pengetahuan Alam*, 13(1): 46-51
- Roy, S. D., & Sen, D. 2025. Two new records of soil nematoda (Dorylaimida: Qudsianematidae) from Odisha, India. *Records of the Zoological Survey of India*, 124(1): 125–132. <https://doi.org/10.26515/rzsi/v124/i1S/2024/172723>
- Rusconi, J. M., García, R., Salas, A., & Achinelly, M. F. 2023. Diversidad de nematodos asociados a Heterodermia diademata: Un estudio de caso en la Eco Área de Avellaneda, Argentina. *Ecología Austral*, 33(3): 798–807. <https://doi.org/10.25260/EA.23.33.3.0.2227>
- Sagita, L., Siswanto, B., dan Kurniatun, H. 2014. Studi keragaman dan kepadatan nematoda pada berbagai sistem penggunaan lahan di sub das konto. *Jurnal Tanah dan Sumberdaya Lahan*.1(1): 51-60.
- Sánchez, M. S. & Ferris, H. 2018. Nematode ecology and soil health. In *Plant parasitic nematodes in subtropical and tropical agriculture* (pp. 62-86). Wallingford UK: CAB International.
- Sarmah, S. 2022. Komunitas nematoda pada lahan pertanaman wortel dan hubungannya dengan populasi mikrob tanah. *Jurnal Tanah dan Iklim*, 46(1): 91-102.
- Shah, M. M., & Mahamood, M. 2017. *Introductory chapter: nematodes-a lesser known group of organisms. Nematology-Concepts, Diagnosis and Control*. InTech. Rijeka. doi: 10.5772/intechopen.68589.
- Sharma, G.C., Gupta, R., 2015. Effect of edaphic factors on Mononchids in the rhizosphere of polyhouse grown flower crops. *Indian J. Nematol.* 45 (1): 12–15.
- Shaw, E. A., Adams, B. J., Barrett, J. E., Lyons, W. B., Virginia, R. A., & Wall, D. H. 2018. Stable C and N isotope ratios reveal soil food web structure and identify the nematode *Eudorylaimus antarcticus* as an omnivore–predator

- in Taylor Valley, Antarctica. *Polar Biology*, 41: 1013-1018.
- Shaw, E. A., Boot, C. M., Moore, J. C., Wall, D. H., & Baron, J. S. 2019. Long-term nitrogen addition shifts the soil nematode community to bacterivore-dominated and reduces its ecological maturity in a subalpine forest. *Soil Biology and Biochemistry*, 130 (1): 177-184.
- Shoaib, M., Mumtaz, S., Abbas, A., Begum, F., & Hussain, B. 2020. 12. Population density of *free-living* nematodes and their relationships with some soil physicochemical properties in forest and agriculture soils of Gapa Valley Chalt, Gilgit, Pakistan. *Pure and Applied Biology (PAB)*, 10(2): 465-473.
- Sidhu, H. S., & Kanwar, R. S. 2020. Effect of prey density of *Meloidogyne incognita* on the predation efficiency of the predator, *fictor composticola* (nematoda: diplogasteridae). *Egyptian Journal of Biological Pest Control*, 30: 1-6.
- Sikder, M. M., & Vestergård, M. 2020. Impacts of root metabolites on soil nematodes. *Frontiers in plant science*, 10 (1): 1792.
- Song, D., Pan, K., Tariq, A., Sun, F., Li, Z., Sun, X., ... & Wu, X. 2017. Large-scale patterns of distribution and diversity of terrestrial nematodes. *Applied Soil Ecology*, 114: 161-169.
- Sopialena, S., Suyadi, S., & Rudiyanto, H. 2022. Keragaman nematoda pada lahan reklamasi pasca tambang batu bara di Desa Bangun Rejo Kecamatan Tenggarong Seberang nematode diversity in post-coal mining reclamation land in Bangun Rejo Village, Tenggarong Seberang District. *Agrifor: Jurnal Ilmu Pertanian dan Kehutanan*, 21(1): 161-174.
- Spaull, V. W. 1973. Distribution of nematode feeding groups at Signy Island, South Orkney Islands, with an estimate of their biomass and oxygen consumption. *British Antarctic Survey Bulletin*, 37 (1): 21-32.
- Sun, X., Zhang X, Zhang S, Dai G, Han S, Liang W. 2013. Soil nematodes responses to increases in nitrogen deposition and precipitation in a temperate forest. *PLoS ONE*, 8 (12): 1-8.
- Suyadi, S., Sila, S., & Samuel, J. 2021. Nematode diversity indices application to determine the soil health status of Lembo agroecosystem in West Kutai, East Kalimantan Province, Indonesia. *Biodiversitas Journal of Biological Diversity*. 22(7): 2861-2869
- Swibawa, I. G. 2001. Keanekaragaman nematoda dalam tanah pada berbagai tipe tataguna lahan di Asb-Benchmark Area Way Kanan. *Jurnal hama dan penyakit tumbuhan tropika*, 1(2): 54-59.
- Syahrok, S. F., & Suryaminarsih, P. 2023. Faktor efektivitas penggunaan nematoda entomopatogen. *Exact Papers in Compilation (EPiC)*, 5(1): 29-33.

- Tahat, M., M. Alananbeh, K., A. Othman, Y., & I. Leskovar, D. 2020. Soil health and sustainable agriculture. *Sustainability*, 12(12): 4859. <https://doi.org/10.3390/su12124859>
- Tahseen, Q., Hussain, A., & Khan, R. 2006. Descriptions of three new and a known species of *Prismatolaimus* de Man, 1880 (Nematoda: Enoplida) from fresh water habitats in India. *J. Nematode Morphol. Syst*, 9 (2): 111-127
- Tarno, H., Marsudi EW, Widjayanti T, Setiawan Y. 2021. Nematodes association with robusta coffee plantations in Malang District, East Java, Indonesia. *Biodiversitas*, 22 (8): 3306-3312.
- Taylor, L. S., Phillips, G., Bernard, E. C., & DeBruyn, J. M. 2020. Soil nematode functional diversity, successional patterns, and indicator taxa associated with vertebrate decomposition hotspots. *PLoS One*, 15(11), e0241777.
- Thoden, T. C., Alkader, M. A., & Wiles, J. A. 2020. Biological attributes of Salibro, a novel sulfonamide nematicide. Part 2: impact on the fitness of various non-target nematodes. *Nematology*, 23(3): 287-303.
- Trap, J., Ranoarisoa, M. P., Raharijaona, S., Rabeharisoa, L., Plassard, C., Mayad, E. H., ... & Blanchart, E. 2021. Agricultural practices modulate the beneficial activity of bacterial-feeding nematodes for plant growth and nutrition: evidence from an original intact soil core technique. *Sustainability*, 13(13): 71-81.
- Traunspurger, W., Reiff, N., Krashevskaya, V., Majdi, N., & Scheu, S. 2017. Diversity and distribution of soil micro-invertebrates across an altitudinal gradient in a tropical montane rainforest of Ecuador, with focus on free-living nematodes. *Pedobiologia*, 62 (1): 28-35
- Tsiafouli, M.A.; Thébault, E.; Sgardelis, S.P.; Ruiter, P.C.; Putten, W.H.; Birkhofer, K.; Hemerik, L.; Vries, F.T.; Bardgett, R.D.; Brady, M.V.; et al. 2015. Intensive agriculture reduces soil biodiversity across Europe. *Glob. Chang. Biol.*, 21: 973–985.
- Vinciguerra, M. T., Eskandari, A., Clausi, M., Asghari, R., & Rappazzo, G. 2016. Description and molecular characterisation of two new species of Dorylaimidae (Dorylaimida: Nematoda) from Iran, with a compendium of all the nominal species of *Dorylaimus* Dujardin, 1845. *Nematology*, 18(3): 293-310.
- Wang, Y. M., Guan, P. T., Chen, J. W., Li, Z. X., Yang, Y. R., & Wang, P. 2021. A comparison of soil nematode community structure and environmental factors along fen-bush-forest succession in a peatland, northeastern China. *Global Ecology and Conservation*, 28(1): e01679.
- Wang, Y., Jiang, H., He, C., Wang, Z., Wang, X., Sheng, L., & Chen, F. 2016. Effect of paddy cultivation on nematode community in Jinchuan Wetland

- of Changbai Mountain area, China. *Ecological Engineering*, 96(1): 20-25.
- Wangge, M. M. N., & Mago, O. Y. T. 2021. Keanekaragaman arthropoda musuh alami hama tanaman kakao (*Theobroma cacao* L.) pada perkebunan polikultur di Desa Hokeng Jaya Kecamatan Wulanggitang Kabupaten Flores Timur. *Spizaetus: Jurnal Biologi Dan Pendidikan Biologi*, 2(1): 47–59.
- Widowati R, Indarti S, Bambang RTP. 2014. Distribution of plant nonparasitic nematode genera on Arabica coffee. *JPTI*, 18 (1): 24-32.
- Wilecki, M., Lightfoot, JW., Susoy, V., and Sommer, RJ. 2015. Predatory feeding behaviour in *Pristionchus* nematodes is dependent on phenotypic plasticity and induced by serotonin. *Journal of Experimental Biology*, 218(9): 1306–1313. doi: 10.1242/jeb.118620.
- Wiyono, W., Hartoyo, E., & Mahananto, M. 2015. Efisiensi pemakaian pupuk urin manusia pada pertumbuhan dan hasil padi sawah. *Jurnal Ilmiah Agrineca*, 15(1).
- Wu, W. J., Xu, C. L., Wang, D. W., & Xie, H. 2018. A new species of the genus *Eudorylaimus* Andr ssy, 1959 (Nematoda: Dorylaimida: Qudsianematidae) associated with *Picea crassifolia* in China. *Zootaxa*, 4526(4): 576-588.
- Wuli, R. N., Loda, W., & Noy, J. A. 2023. Pengaruh jarak tanam pada sistem jajar legowo terhadap produktivitas padi Varietas Inpari 30 di Desa Pape Kecamatan Bajawa Kabupaten Ngada. *Jurnal Pertanian Unggul*, 2(2): 1-9.
- Yadav, S., Patil, J., & Kanwar, R. S. 2018. The role of free living nematode population in the organic matter recycling. *International Journal of Current Microbiology and Applied Sciences*, 7(6): 1-9.
- Yeates, G.W., T. Bongers, R. G. M. De Goede, D. W. Freckman, & S. S. Georgieva. 1993. Feeding habits in soil nematode families and genera—An outline for soil ecologists. *Journal of Nematology*, 25(1): 315-331
- Yusrianti, R., & Japa, L. 2024. Diversity of macrozoobenthos as bioindicator for water quality of Jangkok River Lombok Island. *Jurnal Biologi Tropis*, 24(2): 9-16.
- Zaini, A. H., & Saitama, A. 2023. Analisa perubahan iklim dan pengaruhnya pada produktivitas tanaman padi di Kabupaten Malang. *Plantropica: Journal of Agricultural Science*, 8(2): 173-180.
- Zhang, Y., Ji, L., & Yang, L. 2021. Abundance and diversity of soil nematode community at different altitudes in cold-temperate montane forests in northeast China. *Global Ecology and Conservation*, 29 (01717): 1-12.

- Zhang, Y., Li, S., Li, H., Wang, R., Zhang, K. Q., & Xu, J. 2020. Fungi-nematode interactions: diversity, ecology, and biocontrol prospects in agriculture. *Journal of fungi (Basel, Switzerland)*, 6(4): 206. <https://doi.org/10.3390/jof6040206>
- Zhao, Z. Q. 2009. New Zealand species of the genus *Tripyla* Bastian, 1865 (Nematoda: Triplonchida: Tripylidae). I: A new species, a new record and key to long-tailed species. *Zootaxa*, 2291(1): 35-50.
- Zhao, Z. Q., & Buckley, T. R. 2009. Phylogenetic analysis of nematode nuclear 18S rDNA sequences indicates the genus *Tripylina* Brzeski, 1963 (Nematoda: Tripylidae de Man, 1876) should be placed in Enoplida. *Zootaxa*, 2238(1): 25-32.
- Zhou, G., Zhang, W., Chen, A., He, M., & Ma, X. 2019. Rapid detection of rice disease based on FCM-KM and faster R-CNN fusion. *IEEE access*, 7(1): 143190-143206.
- Zhu, T., Yang, C., Wang, J., Zeng, S., Liu, M., Yang, J., ... & Müller, C. 2018. Bacterivore nematodes stimulate soil gross N transformation rates depending on their species. *Biology and Fertility of Soils*, 54 (1): 107-118.

