

DAFTAR PUSTAKA

- Agus, N., Dian, B., Elly Tuti, W., Aswi Andreasary, M. (2023). First Record of Red Sea Bream, *Pagrus major* in the Eastern Indian Ocean South of Java, Indonesia Revealed by DNA Barcoding. *Biodiversitas*, 24(11), 6023-6030.
- Aisyah, S., Fahrul Syarif, A., Indrawati, A., Manajemen Sumberdaya Perairan, J., Pertanian Perikanan dan Biologi, F., Identifikasi Ikan Selangkat Berdasarkan Karakter Morfologi dan Molekuler di Perairan Kabupaten Bangka Selatan. *Indonesian Journal of Fisheries Science and Technology*, 18(2), 1-6.
- Amal, K., Biswas, Manabu, S., Yoshimasa, T., Kenji, T., Hidemi, K. (2006). Effect of Photoperiod Manipulation on the Growth Performance and Stress Response of Juvenile Red Sea Bream (*Pagrus major*). *Elsevier Aquaculture*, 258, 350-356.
- Artika, R., Sigit, R., Ajiriyanto, M. K., & Yulianto, T. (2022). Image Enhancement Citra Digital Short PIN PWR FUEL Berisi Pelet UO2 Alam Pasca Uji Radiografi Sinar-X Menggunakan Program ImageJ. *Urania : Jurnal Ilmiah Daur Bahan Bakar Nuklir*, 28(1), 21.
- Bintoro, W., Purwanto, E., & Indah Noviyani Jurusan Psikologi, D. (2013). Hubungan Self Regulated Learning dengan Kecurangan Akademik Mahasiswa. *Educational Psychology Journal*, 2(1), 57-64.
- Fadhil, R., Muchlisin, Z. A., Sari Fakultas Kelautan dan Perikanan Universitas Syiah Kuala, W., Aceh, B., Hubungan Panjang-Berat dan Morfometrik Ikan Julung-Julung (*Zenarchopterus dispar*) dari Perairan Pantai Utara Aceh. *Jurnal Ilmiah Mahasiswa Kelautan dan Perikanan Unsyiah*, 1, 146-159.
- Fukuhara, O. (1985). Functional Morphology and Behavior of Early Life Stages of Red Sea Bream. *Japanese Society of Scientific Fisheries*, 51(5), 731-743.
- Google Maps, 2025. Maruki Suisan Fisheries Co.Ltd. <https://www.google.com/maps/place/Amakusa,+Prefektur+Kumamoto,+Jepang/>. Diakses pada 17 April 2025, dari Google Maps: <https://www.google.co.id/maps>.
- Gonzalez, E. B., Aritaki, M., Knutsen, H., & Taniguchi, N. (2015). Effects of Large-Scale Releases on the Genetic Structure of Red Sea Bream (*Pagrus major*, Temminck et Schlegel) Populations in Japan. *PLOS ONE*, 10(5), 1-20.
- Gunathilaka, B. E., Jeong, S. M., Cho, M. U., Kim, K. W., Hur, S. W., Lee, S., You, S. G., & Lee, S. M. (2023). Effects of Dietary Fish Meal Replacement with Alternative Protein Ingredients and Their Combinations on Growth Performance, Feed Utilization, Fillet Composition, and Biochemical

- Parameters of Red Sea Bream (*Pagrus major*). *Aquaculture Nutrition*, 2023, 1–16.
- Islamiyah, S. Al. (2021). Kesesuaian Mutu Ikan Roa (*Hemirhamphus sp.*) Asap dengan Metode Pengasapan Tradisional dan Metode Asap Cair. *Jurnal Sains Dan Teknologi Hasil Pertanian*, 1(2), 53–63.
- Kadim, M. K., Antonio Sinulingga, L., Pasingi, N., & Ibrahim Miolo, M. (2024). Morfometrik dan Meristik Ikan Gobi di Sungai Bone, Gorontalo. *Jurnal Sumberdaya Akuatik Indopasifik*, 8(2), 165–172.
- Karama, K. S., Matsushita, Y., Inoue, M., Kojima, K., Tone, K., Nakamura, I., & Kawabe, R. (2021). Movement Pattern of Red Sea Bream (*Pagrus major*) and Yellowtail (*Seriola Quinqueradiata*) around Offshore Wind Turbine and the Neighboring Habitats in the Waters near Goto Islands, Japan. *Aquaculture and Fisheries*, 6(3), 300–308.
- Kato, K. (2023). Breeding Studies on Red Sea Bream (*Pagrus major*): Mass Selection to Genome Editing. *Fisheries Science*, 89(2), 103–119.
- Koshio, S. (2002). Red Sea Bream, *Pagrus major*. *Nutrient Requirements*, 4(2), 51–63.
- Mana, R. R., & Kawamura, G. (2002). A Comparative Study on Morphological Differences in the Olfactory System of Red Sea Bream (*Pagrus major*) and Black Sea Bream (*Acanthopagrus schlegeli*) from Wild & Cultured Stocks. *Elsevier Aquaculture*, 209, 285–306.
- Matthews, W. J. (1998). Patterns in Freshwater Fish Ecology. In Patterns in Freshwater Fish Ecology (Vol. 2). Springer US.
- Natural Atlas. (2005). Red Sea Bream. <https://naturalatlas.com/>
- Primawati, S. N., Efendi, I., & Marnita. (2016). Identifikasi Jenis Ikan Hasil Tangkapan Nelayan di Pantai Jeranjang. *Jurnal Pendidikan Mandala*, 1, 73–78.
- Rismanto, M., Gustomi¹, A., & Adibrata, S. (2023). Karakteristik Morfometrik & Meristik Ikan Gabus (*Channa striata*) pada Beberapa Tipe Perairan di Pulau Bangka. *Jurnal Sumberdaya Perairan*, 17(1), 12–18.
- Saroni, S., Al Bohari, I., Darwa Suryani, S., Biologi, P., & Muhammadiyah Bengkulu, U. (2024). Perbandingan Morfometrik Ikan Nila (*Oreochromis niloticus*) Budidaya dengan Sungai di Air Kedurang Kecamatan Kedurang Kabupaten Bengkulu Selatan. *Jurnal Riset Dan Inovasi Pendidikan Sains*, 3(2), 109–117.
- Takács, P., Vitál, Z., Ferincz, Á., & Staszny, Á. (2016). Repeatability, Reproducibility, Separative Power and Subjectivity of Different Fish Morphometric Analysis Methods. *PLOS ONE*, 11(6), 1–16.

- Takahashi, K., & Masuda, R. (2019). Nurture is above Nature: Nursery Experience Determines Habitat Preference of Red Sea Bream (*Pagrus major*) Juveniles. *Journal of Ethology*, 37(3), 317–323.
- Yamaguchi, T., Takayama, K., Hirose, N., & Matsuyama, M. (2020). The Sea of Amakusa Playing the Role of a Distributor of Swordtip Squid (*Uroteuthis edulis*) Migrating from the East China Sea to the East and West Sides of Japan. *Fisheries Research*, 225(20), 1–7.
- Yamaha. (1990). The “King of Fish”, its History and Development as a fishery. *Fishery Journal*, 33(3), 1–8.
- Yasuda, T., Komeyama, K., Kato, K., Mitsunaga, Y. (2012). Use of Acceleration Loggers in Aquaculture to Determine Net-Cage Use and Fiel Metabolic Rates in Red Sea Bream *Pagrus major*. *The Japanese Society of Fisheries Science*, 78:229-235

