

REFERENCES

- Bennett, L.E., Jegasothy, H., Konczak, I., Frank, D., Sudharmarajan, S. & Clingeffer, P.R., 2011. Total polyphenolics and anti-oxidant properties of selected dried fruits and relationships to drying conditions. *Journal of functional foods*, 3(2), pp.115-124.
- Bernard, D., Kwabena, A.I., Osei, O.D., Daniel, G.A., Elom, S.A. & Sandra, A., 2014. The effect of different drying methods on the phytochemicals and radical scavenging activity of Ceylon Cinnamon (*Cinnamomum zeylanicum*) plant parts. *European Journal of Medicinal Plants*, 4(11), pp.1324-1335.
- Brandle, J.E., Starratt, A.N. & Gijzen, M., 1998. Stevia rebaudiana: Its agricultural, biological, and chemical properties. *Canadian Journal of Plant Science*, 78(4), pp.527-536.
- Ceunen, S. & Geuns, J.M., 2013. Steviol glycosides: chemical diversity, metabolism, and function. *Journal of Natural Products*, 76(6), pp.1201-1228.
- Chandra, A. & Novalia, N., 2014. Studi awal ekstraksi batch daun *Stevia rebaudiana* Bertoni dengan variabel jenis pelarut dan temperatur. *Research Report-Engineering Science*, 2, pp.1-8.
- Chew, K.K., Khoo, M.Z., Ng, S.Y., Thoo, Y.Y., Aida, W.W. & Ho, C.W., 2011. Effect of ethanol concentration, extraction time and extraction temperature on the recovery of phenolic compounds and antioxidant capacity of *Orthosiphon stamineus* extracts. *International Food Research Journal*, 18(4), p.1427.
- Chu, D.C. & Juneja, L.R., 1997. General chemical composition of green tea and its infusion. *Chemistry and applications of green tea*, 2(1), pp.13-22.
- Chun, O.K., Kim, D.O. & Lee, C.Y., 2003. Superoxide radical scavenging activity of the major polyphenols in fresh plums. *Journal of Agricultural and Food Chemistry*, 51(27), pp.8067-8072.
- Darmawan, W. & Mahayasih, P.G.M.W., 2023. Pengaruh metode pengeringan terhadap aktivitas antioksidan ekstrak *Stevia rebaudiana* bertoni melalui ekstraksi berbantuan gelombang mikro. *An-Najat*, 1(3), pp.117-128.
- Fatimah, S., 2022. Kandungan Metabolit Sekunder Daun Muda dan Daun Dewasa Tumbuhan Keji (*Staurogyne elongata* [Blume] Kuntze) di Kabupaten Pekalongan (Doctoral dissertation, Thesis, Universitas Islam Negeri Walisongo Semarang).
- Goyal, S.K., Samsher, N. & Goyal, R., 2010. Stevia (*Stevia rebaudiana*) a bio-sweetener: a review. *International Journal of Food Sciences and Nutrition*, 61(1), pp.1-10.
- Halim, A. A., Zain, Z. M., Mubarak, A., & Ahmad, F. T., 2019. Effect of different drying methods on antioxidant properties, stevioside, and rebaudioside A contents of stevia (*Stevia rebaudiana* Bertoni) leaves. *Asian Journal of Agriculture and Biology*, 7(1), pp.61-68.

- Handayani, D., Mun'im, A. & Ranti, A.S., 2014. Optimation of green tea waste axtraction using microwave assisted extraction to yield green tea extract. *Majalah Obat Tradisional*, 19(1), pp.29-35.
- Hossain, M.N., Sarker, U., Raihan, M.S., Al-Huqail, A.A., Siddiqui, M.H. & Oba, S., 2022. Influence of salinity stress on color parameters, leaf pigmentation, polyphenol and flavonoid contents, and antioxidant activity of *Amaranthus lividus* leafy vegetables. *Molecules*, 27(6), p.1821.
- Lemus-Mondaca, R., Vega-Gálvez, A., Zura-Bravo, L., & Ah-Hen, K., 2012. *Stevia rebaudiana* Bertoni, source of a high-potency natural sweetener: A comprehensive review on the biochemical, nutritional and functional aspects. *Food chemistry*, 132 3, pp.1121-1132.
- Madan, S., S. Ahmad, G. N. Singh, K. Kohli, Y. Kumar, R. Singh. & M. Garg. 2010. *Stevia rebaudiana* (Bert) Bertoni. *Indian Journal of Natural Products and Resources*, 3, pp.267-280.
- Mokrani, A. & Madani, K., 2016. Effect of solvent, time and temperature on the extraction of phenolic compounds and antioxidant capacity of peach (*Prunus persica* L.) fruit. *Separation and Purification Technology*, 162, pp.68-76.
- Müller, J. & Heindl, A., 2006. Drying of medicinal plants. In: R.J. Bogers, L.E. Craker, and D. Lange, eds. *Medicinal and Aromatic Plants*. Dordrecht: Springer, pp.237-252.
- Myint, K.Z., Zhou, Z., Shi, Q., Chen, J., Dong, X. & Xia, Y., 2023. *Stevia* polyphenols, their antimicrobial and anti-inflammatory properties, and inhibitory effect on digestive enzymes. *Molecules*, 28(22), p.7572.
- Rahmawati, M. 2018. Pengaruh Kadar Polifenol terhadap Warna Daun Tembakau Vorstenlenden (*Nicotiana tabacum* var. vorstenlenden) di PT. Perkebunan Nusantara X (Skripsi, Universitas Gadjah Mada).
- Rukmana, I.H.R., 2003. *Budi Daya Stevia, Bahan Pembuatan Pemanis Alami*. Kanisius.
- Siddiq, H.B.H.F., Riyuwani, J. & Dewi, R.D.Y., 2017. Penentuan kadar polifenol ekstrak teh kemasan dengan metode remaserasi menggunakan spektrofotometri UV-Vis. *Jurnal Ilmiah Farmasi Akademi Farmasi Jember*, 2(1), pp.7-14.
- Singleton, V.L., Orthofer, R., & Lamuela-Raventós, R.M., 1999. Analysis of total phenols and other oxidation substrates and antioxidants by means of Folin-Ciocalteu reagent. *Methods in Enzymology*, 299, pp.152-178.
- Stalikas, C.D., 2007. Extraction, separation, and detection methods for phenolic acids and flavonoids. *Journal of Separation Science*, 30(18), pp.3268-3295.

- Susanti, D.Y. 2008. Efek Suhu Pengeringan terhadap kandungan fenolik dan kandungan katekin ekstrak daun kering gambir. *Prosiding Seminar Nasional Teknik Pertanian Universitas Gadjah Mada Yogyakarta*, 208, pp.123-130.
- Tuminah, S., 2004. Teh *Camellia sinensis* OK Var. Assamica (mast) sebagai salah satu sumber antioksidan. *Cermin dunia kedokteran*, 144, pp.52-54.
- Widarta, I.W.R. & Arnata, I.W., 2017. Ekstraksi komponen bioaktif daun alpukat dengan bantuan ultrasonik pada berbagai jenis dan konsentrasi pelarut. *Agritech*, 37(2), pp.148-157.

