

DAFTAR PUSTAKA

- Alongi, D. M. (2020). Carbon cycling in the world's mangrove ecosystems revisited: Significant sinks and sources. In *Blue Carbon Ecosystems for Sustainable Development* (pp. 31–74). Springer. https://doi.org/10.1007/978-981-15-2164-4_3
- Ardli, E. R., & Wolff, M. (2008). Quantifying habitat and resource use changes in the Segara Anakan Lagoon (Cilacap, Indonesia) over the past 25 years (1978–2004). *Asian Journal of Water, Environment and Pollution*, 5(4), 59–67.
- Ardli, E. R., & Wolff, M. (2009). Land use and land cover change affecting habitat distribution in the Segara Anakan Lagoon, Java, Indonesia. *Regional Environmental Change*, 9(4), 235–243. <https://doi.org/10.1007/s10113-008-0072-6>
- Chave, J., Andalo, C., Brown, S., Cairns, M. A., Chambers, J. Q., Eamus, D., Fölster, H., Fromard, F., Higuchi, N., Kira, T., Lescure, J.-P., Nelson, B. W., Ogawa, H., Puig, H., Riéra, B., & Yamakura, T. (2005). Tree allometry and improved estimation of carbon stocks and balance in tropical forests. *Oecologia*, 145(1), 87–99. <https://doi.org/10.1007/s00442-005-0100-x>
- Clough, B. F., & Scott, K. (1989). Allometric relationships for estimating above-ground biomass in six mangrove species. *Forest Ecology and Management*, 27(2), 117–127. [https://doi.org/10.1016/0378-1127\(89\)90034-0](https://doi.org/10.1016/0378-1127(89)90034-0)
- Dalponte, M., & Coomes, D. A. (2016). Tree-centric mapping of forest carbon density from airborne laser scanning and hyperspectral data. *Methods in Ecology and Evolution*, 7(10), 1236–1245. <https://doi.org/10.1111/2041-210X.12575>
- Donato, D. C., Kauffman, J. B., Murdiyarso, D., Kurnianto, S., Stidham, M., & Kanninen, M. (2011). Mangroves among the most carbon-rich forests in the tropics. *Nature Geoscience*, 4(5), 293–297. <https://doi.org/10.1038/ngeo1123>
- Dong, P., & Chen, Q. (2017). *LiDAR remote sensing and applications*. CRC Press.
- Duke, N. C., Meynecke, J. O., Dittmann, S., Ellison, A. M., Anger, K., Berger, U., Cannicci, S., Diele, K., Ewel, K. C., Field, C. D., Koedam, N., Lee, S. Y., Marchand, C., Nordhaus, I., & Dahdouh-Guebas, F. (2007). A world without mangroves? *Science*, 317(5834), 41–42. <https://doi.org/10.1126/science.317.5834.41b>
- Eddy, S., Ridho, M. R., Iskandar, I., & Mulyana, A. (2016). Community-based mangrove forest conservation for sustainable fisheries. *Jurnal Silvikultur Tropika*, 7(3), 42–47.
- Fatoyinbo, T. E., & Simard, M. (2013). Height and biomass of mangroves in Africa from ICESat/GLAS and SRTM. *International Journal of Remote Sensing*, 34(2), 668–681. <https://doi.org/10.1080/01431161.2012.712224>
- Fatoyinbo, T., Lagomasino, D., Feliciano, E. A., Lee, S. K., Trettin, C. C., & Simard, M. (2018). Estimating mangrove aboveground biomass from airborne LiDAR data: A case study from the Zambezi River delta. *Environmental Research Letters*, 13(2), 025012. <https://doi.org/10.1088/1748-9326/aa9f03>

- Feliciano, E. A., Wdowinski, S., Potts, M. D., Lee, S. K., & Fatoyinbo, T. E. (2017). Estimating mangrove canopy height and above-ground biomass in the Everglades National Park with airborne LiDAR and TanDEM-X data. *Remote Sensing*, 9(7), 702. <https://doi.org/10.3390/rs9070702>
- Fromard, F., Puig, H., Mougin, E., Marty, G., Betoulle, J. L., & Cadamuro, L. (1998). Structure, above-ground biomass and dynamics of mangrove ecosystems: New data from French Guiana. *Oecologia*, 115(1–2), 39–53. <https://doi.org/10.1007/s004420050489>
- Getzner, M., & Islam, M. S. (2020). Ecosystem services of mangrove forests: Results of a meta-analysis of economic values. *International Journal of Environmental Research and Public Health*, 17(16), 5830. <https://doi.org/10.3390/ijerph17165830>
- Giri, C., Ochieng, E., Tieszen, L. L., Zhu, Z., Singh, A., Loveland, T., Duke, N. (2011). Status and distribution of mangrove forests of the world using earth observation satellite data. *Global Ecology and Biogeography*, 20(1), 154–159. <https://doi.org/10.1111/j.1466-8238.2010.00584.x>
- Holtermann, T., Gade, M., Lange, P. K., & Körtzinger, A. (2009). Remote sensing of suspended particulate matter in the Segara Anakan Lagoon, Java, Indonesia. *Estuarine, Coastal and Shelf Science*, 83(4), 459–471. <https://doi.org/10.1016/j.ecss.2009.04.032>
- Kauffman, J. B., & Donato, D. C. (2012). *Protocols for the measurement, monitoring, and reporting of structure, biomass, and carbon stocks in mangrove forests*. CIFOR.
- Kathiresan, K., & Bingham, B. L. (2001). Biology of mangroves and mangrove ecosystems. *Advances in Marine Biology*, 40, 81–251. [https://doi.org/10.1016/S0065-2881\(01\)40003-4](https://doi.org/10.1016/S0065-2881(01)40003-4)
- Komiyama, A., Ong, J. E., & Pongparn, S. (2008). Allometry, biomass, and productivity of mangrove forests: A review. *Aquatic Botany*, 89(2), 128–137. <https://doi.org/10.1016/j.aquabot.2007.12.006>
- Marcello, J., Spinola, M., Albors, L., Marques, F., Rodríguez-Esparragón, D., & Eugenio, F. (2024). Performance of individual tree segmentation algorithms in forest ecosystems using UAV LiDAR data. *Drones*, 8(12), 772. <https://doi.org/10.3390/drones8120772>
- Nemmaoui, A., Aguilar, F. J., & Aguilar, M. A. (2024). Benchmarking of individual tree segmentation methods in Mediterranean forest based on point clouds from UAV imagery and low-density airborne laser scanning. *Remote Sensing*, 16(21), 3974. <https://doi.org/10.3390/rs16213974>
- Owers, C. J., Rogers, K., & Saintilan, N. (2018). Allometric equations for aboveground biomass in mangrove species of Southeast Australia. *Wetlands Ecology and Management*, 26(4), 563–575. <https://doi.org/10.1007/s11273-018-9584-5>
- Pham, T. D., et al. (2019). Aboveground biomass estimation of mangrove forests using canopy height and other LiDAR metrics. *Remote Sensing*, 11(15), 1780. <https://doi.org/10.3390/rs11151780>

- Ploton, P., Barbier, N., Takoudjou, M. D., et al. (2016). Closing a gap in tropical forest biomass estimation: Widespread prediction of canopy height from crown diameter measurements. *Biogeosciences*, 13(5), 1571–1585. <https://doi.org/10.5194/bg-13-1571-2016>
- Portet, S. (2020). A primer on model selection using the Akaike information criterion. *Infectious Disease Modelling*, 5, 111–128. <https://doi.org/10.1016/j.idm.2019.12.010>
- Salum, R. B., Souza-Filho, P. W. M., Simard, M., Silva, C. A., Fernandes, M. E. B., Cougo, M. F., do Nascimento, W., & Rogers, K. (2020). Improving mangrove above-ground biomass estimates using LiDAR. *Estuarine, Coastal and Shelf Science*, 236, 106585. <https://doi.org/10.1016/j.ecss.2020.106585>
- Suwa, R., et al. (2020). A novel approach to estimating above-ground biomass of mangroves using tree height alone. *Forests*, 11(3), 345. <https://doi.org/10.3390/f11030345>
- Tang, H., Brolly, M., Zhao, F., et al. (2018). LiDAR remote sensing for vegetation structural parameters: A review. *ISPRS Journal of Photogrammetry and Remote Sensing*, 144, 1–17. <https://doi.org/10.1016/j.isprsjprs.2018.06.003>
- Zhang, Z., et al. (2023). Model selection and evaluation in ecological studies: Balancing fit and complexity with AIC. *Ecological Modelling*, 476, 110216. <https://doi.org/10.1016/j.ecolmodel.2023.110216>

