

DAFTAR PUSTAKA

- Allen. (2001). Report on the Present State of Knowledge with regard to the Habits and Migrations of the Mackerel (*Scomber scomber*). *Marine Biological Association*, 5 (1).
- Araújo, M. S., Perez, S. I., Magazoni, M. J. C., & Petry, A. C. (2014). Body size and allometric shape variation in the molly *Poecilia vivipara* along a gradient of salinity and predation. *BMC Evolutionary Biology*, 14(1). <https://doi.org/10.1186/s12862-014-0251-7>
- Cai, K., Kindong, R., Ma, Q., Han, X., & Qin, S. (2022). North Pacific Fisheries Commission Chub mackerel (*Scomber japonicus*). *Marine Science and Engineering*.
- Caves, E. M., Sutton, T. T., & Johnsen, S. (2017). Visual acuity in ray-finned fishes correlates with eye size and habitat. *Journal of Experimental Biology*, 220(9), 1586–1596. <https://doi.org/10.1242/jeb.151183>
- Čikeš, K., & Barbara, Z. (2011). Biometry Markers of Chub Mackerel, *Scomber japonicus* Houttuyn, 1782, in the Adriatic Sea. *Acta Adriatica*.
- Collette, B. (1983). Scombrids of the World: an Annotated and Illustrated Catalogue of Tunas, Mackerels, Bonitos, and Related Species Known to Date. *FAO Fisheries Synopsis*.
- Erguden, D., Öztürk, B., Aka Erdogan, Z., & Turan, C. (2009). Morphologic structuring between populations of chub mackerel *Scomber japonicus* in the Black, Marmara, Aegean, and northeastern Mediterranean Seas. *Fisheries Science*, 75(1), 129–135. <https://doi.org/10.1007/s12562-008-0032-6>
- Friedman, S. T., Price, S. A., & Wainwright, P. C. (2021). The Effect of Locomotion Mode on Body Shape Evolution in Teleost Fishes. *Integrative Organismal Biology*, 3(1). <https://doi.org/10.1093/iob/obab016>
- Froese, & Pauly. (2024). *Scomber japonicus* Houttuyn, 1782 Chub mackerel.
- Geladakis, G., Nikolioudakis, N., Koumoundouros, G., & Somarakis, S. (2018). Morphometric discrimination of pelagic fish stocks challenged by variation in body condition. *ICES Journal of Marine Science*, 75(2), 711–718. <https://doi.org/10.1093/icesjms/fsx186>
- Hart. (1973). *Scomber japonicus*. *Fisheries Research Board of Canada*.
- Hernández, J. (2000). Synopsis of Biological Data on the Chub Mackerel (*Scomber japonicus* Houttuyn, 1782). *FAO Fisheries Synopsis*.
- Hernandez, J., Villalobos-Leiva, A., Bermúdez, A., Ahumada-C, D., Suazo, M. J., Correa, M., Díaz, A., & Benítez, H. A. (2022). Ecomorphology and Morphological Disparity of *Caquetaia kraussii* (Perciformes: Cichlidae) in Colombia. *Animals*, 12(23). <https://doi.org/10.3390/ani12233438>

- Higham, T. E. (2007). The integration of locomotion and prey capture in vertebrates: Morphology, behavior, and performance. *Integrative and Comparative Biology*, 47(1), 82–95. <https://doi.org/10.1093/icb/icm021>
- Ihssen, P. E., Booke, H. E., Casselman, J. M., McGlade, J. M., Payne, N. R., & Utter, F. M. (1981). Stock Identification: Materials and Methods. *Canadian Journal of Fisheries and Aquatic Sciences*, 38(12), 1838–1855. <https://doi.org/10.1139/f81-230>
- Jannat, B., Kanon, K. F., Sultana, N., Alam, M. S., & Alam, M. S. (2022). Morphological Variations Between Native and Vietnam-originated striped snakeheads (*Channa striata*) in Bangladesh. *Bangladesh Journal of Fisheries*, 34(1), 41–52. <https://doi.org/10.52168/bjf.2022.34.5>
- Keč, V. (2011). Biometry Markers of Chub Mackerel, *Scomber japonicus* Houttuyn, 1782, in the Adriatic Sea. *Institute of Oceanography and Fisheries*, 52(2).
- Lee, D., Son, S. H., Kim, W., Park, J. M., Joo, H., & Lee, S. H. (2018). Spatio-Temporal Variability of the Habitat Suitability Index for Chub Mackerel (*Scomber japonicus*) in the East/Japan Sea and the South Sea of South Korea. *Remote Sensing*, 10(6). <https://doi.org/10.3390/rs10060938>
- Lisney, T. J., Collin, S. P., & Kelley, J. L. (2020). The effect of ecological factors on eye morphology in the western rainbowfish, *Melanotaenia australis*. *Journal of Experimental Biology*, 223(10). <https://doi.org/10.1242/jeb.223644>
- Maruki, T. (2014). Geography of Amakusa City, Kumamoto Prefecture, Japan. *Amakusa Guide*.
- Mashar, A., Wahyuni, Y. S., Hakim, A. A., Wardiatno, Y., Manajemen, D., Perairan, S., Perikanan, F., Kelautan, I., & Pertanian Bogor, I. (2019). Pendekatan Truss Morphometric Dalam Menganalisis Kekerabatan Populasi *Cherax quadricarinatus* (Von Martens, 1868) Di Perairan Jawa Barat (Truss Morphometric Approach for Population Kinship Analysis of *Cherax quadricarinatus* (Von Martens, 1868) in West Java Waters). *Journal of Tropical Fisheries Management*, 3 (2).
- Muniz, A. A., Moura, A., Triay-Portella, R., Moreira, C., Santos, P. T., & Correia, A. T. (2021). Population structure of the chub mackerel (*Scomber colias*) in the North-east Atlantic inferred from otolith shape and body morphometrics. *Marine and Freshwater Research*, 72(3), 341–352. <https://doi.org/10.1071/MF19389>
- Murta, A. G. (2000). Morphological variation of horse mackerel (*Trachurus trachurus*) in the Iberian and North African Atlantic: Implications for stock identification. *ICES Journal of Marine Science*, 57(4), 1240–1248. <https://doi.org/10.1006/jmsc.2000.0810>

- Nauen, J. C., & Lauder, G. V. (2002). Locomotion in scombrid fishes: morphology and kinematics of the finlets of the chub mackerel (*Scomber japonicus*). *Journal of Experimental Biology*, 205 (15), 2143–2154.
- Oktavia, S., Setiawan, U., Nurpadiana, H., & History, A. (2020). Morphological Character Analysis of Mackerel (*Scomberomorus commerson* Lac., 1800) in Sunda Strait. *BIOSFER: Jurnal Tadris Biologi*, 11(1), 1–9. <https://doi.org/10.24042/biosfer>
- Oufiero, C. E., & Whitlow, K. R. (2016). The evolution of phenotypic plasticity in fish swimming. *Current Zoology*, 62(5), 475–488. <https://doi.org/10.1093/cz/zow084>
- Sarower, E. M. M., Khatun, A., Boidya, P., & Samad, M. A. (2019). Meristic and morphometric variations of barred spiny eel *Macrognahtus pancalus* populations from bangladeshi freshwaters: An insight into landmark-based truss network system. *Croatian Journal of Fisheries*, 77(1), 7–18. <https://doi.org/10.2478/cjf-2019-0002>
- Sarower-E-Mahfuj, M., Khatun, A., Boidya, P., & Samad, M. A. (2019). Meristic and morphometric variations of barred spiny eel *Macrognahtus pancalus* populations from bangladeshi freshwaters: An insight into landmark-based truss network system. *Ribarstvo, Croatian Journal of Fisheries*, 77(1), 7–18. <https://doi.org/10.2478/cjf-2019-0002>
- Schakmann, M., & Korsmeyer, K. E. (2023). Fish swimming mode and body morphology affect the energetics of swimming in a wave-surge water flow. *Journal of Experimental Biology*, 226(6). <https://doi.org/10.1242/jeb.244739>
- Sugiono. (2017). Metode Penelitian Kuantitatif dan Kualitatif. *Alfabeta*.
- Turan, C. (1999). A Note on The Examination of Morphometric Differentiation Among Fish Populations: The Truss System. *Tr. J. of Zoology*, 23, 259–263.
- Wang, J., Wainwright, D. K., Lindengren, R. E., Lauder, G. V., & Dong, H. (2020). Tuna locomotion: A computational hydrodynamic analysis of finlet function. *Journal of the Royal Society Interface*, 17(165). <https://doi.org/10.1098/rsif.2019.0590>
- Yoon, S., Kim, D., Baeck, B., & Kim, J. (2008). Feeding Habits of Chub Mackerel (*Scomber japonicus*) in the South Sea of Korea. *J.Kor.Fish.Soc*, 4(1).
- Yukami. (2023). Distribution of Chub mackerel (*Scomber japonicus*). *North Pacific Fisheries Commission*, 8 (15), 1–12. www.npfc.int
- Zhao, G., Zhang, H., & Tang, F. (2025). The Spatio-Temporal Distribution and Population Dynamics of Chub Mackerel (*Scomber japonicus*) in the High Seas of the Northwest Pacific Ocean. *Animals*, 15(8). <https://doi.org/10.3390/ani15081135>